Please check the examination deta	ils below before en	tering your candidate information
Candidate surname		Other names
Pearson Edexcel International Advanced Level	Centre Numbe	r Candidate Number
Time 1 hour 30 minutes	Paper referenc	WME02/01
Mathematics		
International Advance Mechanics M2	d Subsidiaı	ry/Advanced Level
You must have: Mathematical Formulae and Stat	tistical Tables (Y	ellow), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided - there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.
- Good luck with your examination.





Turn over 🕨



1. A motorcyclist and his motorcycle have a combined mass of 480 kg.	
The motorcyclist drives down a straight road that is inclined at an angle θ to the horizontal,	
where $\sin \theta = \frac{1}{12}$, with the engine of the motorcycle working at 3.5 kW. The motorcycle	
is moving at a constant speed of $V \text{m s}^{-1}$. The resistance to the motion of the motorcycle is modelled as a constant force with magnitude 20 <i>V</i> newtons.	
Find the value of V.	
(5)	

Question 1 continued		bl
	(Total 5 marks)	Q
	(iotai 3 mai KS)	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	Tui	rn ov

DO NOT WRITE IN THIS AREA

At time t = 0, *P* is at the origin *O*. (b) Find the position vector of P relative to O at the instant when P is moving in the direction of the vector **j** (4) 4 P 6 5 7 9 4 A 0 4 2 8

A particle P of mass 1.5 kg moves under the action of a single force \mathbf{F} newtons. 2.

At time *t* seconds, $t \ge 0$, *P* has velocity $\mathbf{v} \mathbf{m} \mathbf{s}^{-1}$, where

$$\mathbf{v} = (5t^2 - t^3)\mathbf{i} + (2t^3 - 8t)\mathbf{j}$$

(a) Find **F** when t = 2

(4)

Leave blank

	Q
(Total 8 marks)	

DO NOT WRITE IN THIS AREA

Leave blank



Figure 1

The uniform lamina *ABCD* is a square of side 6a. The template *T*, shown shaded in Figure 1, is formed by removing the right-angled triangle *EFG* and the circle, centre *H* and radius *a*, from the square lamina.

Triangle EFG has EF = EG = 4a, with EF parallel to AB and EG parallel to AD. The distance between AB and EF is a and the distance between AD and EG is a.

The point H lies on AC and the distance of H from BC is 2a.

(a) Show that the centre of mass of *T* is a distance $\frac{4(67 - 3\pi)}{3(28 - \pi)}a$ from *AD*.

The template T is suspended from the ceiling by two light inextensible vertical strings. One string is attached to T at A and the other string is attached to T at B so that T hangs in equilibrium with AB horizontal.

The weight of T is W. The tension in the string attached to T at B is kW, where k is a constant.

(b) Find the value of k, giving your answer to 2 decimal places.

(3)

(5)



3.

Question 3 continued	

DO NOT WRITE IN THIS AREA

lestion 3 continued	Leave blank

Question 3 continued		
		C
	(Total 8 marks)	
		'n o

DO NOT WRITE IN THIS AREA

Leave blank



(6)

DO NOT WRITE IN THIS AREA



A particle *P* of mass 0.3 kg is moving with speed 5 m s^{-1} along a straight line on a smooth horizontal plane. The particle receives a horizontal impulse of magnitude *J*Ns. The speed of *P* immediately after receiving the impulse is 8 m s^{-1} . The angle between the direction of motion of *P* before it receives the impulse and the direction of the impulse is 60° , as shown in Figure 2.

Find the value of *J*.

4.



Question 4 continued		Leav
		Q4
	(Total 6 marks)	
		11
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	Tui	rn over

DO NOT WRITE IN THIS AREA

Leave blank

(3)

(6)

DO NOT WRITE IN THIS AREA



C

The magnitude of the force acting on the rod at A is λMg , where λ is a constant.

(b) Find, to 2 significant figures, the value of λ .

12

5.

Question 5 continued			
		······································	

DO NOT WRITE IN THIS AREA

Question 5 continued	Leave blank
Zutstion 5 continueu	

Question 5 continued		Leav
	(Total 0 1)	Q5
	(Total 9 marks)	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	Tu	15 rn ovei



Figure 4

Two particles, A and B, of mass 2kg and 3kg respectively, are connected by a light inextensible string. Particle A is held at rest at the point X on a fixed rough ramp that is inclined at an angle θ to the horizontal, where $\tan \theta = \frac{5}{12}$. The string passes over a small smooth pulley P that is fixed at the top of the ramp. Particle B hangs vertically below P, 2m above the ground, as shown in Figure 4.

The particles are released from rest with the string taut so that A moves up the ramp and the section of the string from A to P is parallel to a line of greatest slope of the ramp.

The coefficient of friction between A and the ramp is $\frac{3}{6}$

Air resistance is ignored.

- (a) Find the potential energy lost by the system as *A* moves 2 m up the ramp.
- (b) Find the work done against friction as *A* moves 2 m up the ramp.

When B hits the ground, B is brought to rest by the impact and does not rebound and A continues to move up the ramp.

(c) Use the work-energy principle to find the speed of B at the instant before it hits the ground.

(4)

(3)

(4)

Particle A comes to instantaneous rest at the point Y on the ramp, where XY = (2 + d)m.

(d) Use the work-energy principle to find the value of *d*.

(4)



DO NOT WRITE IN THIS ARE

Question 6 continued	

DO NOT WRITE IN THIS AREA

lestion 6 continued	

Question 6 continued	blank	
	Q6	
(Total 15 marks)		
	19]

DO NOT WRITE IN THIS AREA

Leave blank

(4)

(4)

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS ARE

7. [*In this question, the unit vectors* **i** *and* **j** *are in a vertical plane,* **i** *being horizontal and* **j** *being vertically upwards.*]



0

A small ball is projected with velocity $(6\mathbf{i} + 12\mathbf{j}) \text{ m s}^{-1}$ from a fixed point *A* on horizontal ground. The ball hits the ground at the point *B*, as shown in Figure 5. The motion of the ball is modelled as a particle moving freely under gravity.

(a) Find the distance *AB*.

When the height of the ball above the ground is more than *h* metres, the speed of the ball is less than 10 m s^{-1}

(b) Find the smallest possible value of *h*.

When the ball is at the point C on its path, the direction of motion of the ball is perpendicular to the direction of motion of the ball at the instant before it hits the ground at B.

(c) Find, in terms of **i** and **j**, the velocity of the ball when it is at *C*.



DO NOT WRITE IN THIS AREA

	Leave blank
uestion 7 continued	

Question 7 continued		blank
	(Total 11 marks)	Q7
	TUTAL II IIIALKS)	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	Turn	23 over

DO NOT WRITE IN THIS AREA

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8. Particles A, B and C, of masses 2m, m and 3m respectively, lie at rest in a straight line on a smooth horizontal plane with B between A and C. Particle A is projected towards particle B with speed 2u and collides directly with B.

The coefficient of restitution between each pair of particles is e.

- (a) (i) Show that the speed of B immediately after the collision with A is $\frac{4}{3}u(1+e)$
 - (ii) Find the speed of A immediately after the collision with B.

(7)

At the instant when A collides with B, particle C is projected with speed u towards B so that B and C collide directly.

(b) Show that there will be a second collision between A and B.

(6)

Question 8 continued	

DO NOT WRITE IN THIS AREA

		Leave blank
Question 8 continu	ed	

Question 8 continued		ł

DO NOT WRITE IN THIS AREA

uestion 8 continued	Leave blank
	Q8
(Total 13 marks)	
TOTAL FOR PAPER: 75 MARKS END	

