



Pearson
Edexcel

Mark Scheme (Results)

Summer 2024

Pearson Edexcel International Advanced Level
In Statistics S1 (WST01) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2024

Question Paper Log Number P75720RA

Publications Code WST01_01_2406_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)

Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN:

- bod – benefit of doubt
- ft – follow through
 - the symbol \surd will be used for correct ft
- cao – correct answer only
- cso – correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC – special case
- oe – or equivalent (and appropriate)
- d... or dep – dependent
- indep – independent
- dp – decimal places
- sf – significant figures
- * – The answer is printed on the paper or ag- answer given
- \square or d... – The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.

6. If a candidate makes more than one attempt at any question:
 - a) If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - b) If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

Special notes for marking Statistics exams (for AAs only)

- Any correct method should gain credit. If you cannot see how to apply the mark scheme but believe the method to be correct then please send to review.
- For method marks, we generally allow or condone a slip or transcription error if these are seen in an expression. We do not, however, condone or allow these errors in accuracy marks.

Question	Scheme		Marks
1 (a)	$k = 3$		B1 (1)
(b)	$Q_1 = 39 \quad Q_3 = 57$		B1 B1 (2)
(c)	"57" $+1.5 \times ("57" - "39")$ or "39" $-1.5 \times ("57" - "39")$		M1
	84 and 12 therefore only 1 outlier [85]		A1 (2)
(d)			M1
			M1
			M1
			A1 (4)
(e)	A correct difference of the medians with supporting figures e.g. On average Birch [trees grow slightly] taller as the median is larger 48 > 45 oe or A correct difference of the spread with supporting figures e.g. Maple has a greater spread/variation of heights as the range is larger 55 > 48 (excluding outlier) oe e.g. Birch has a greater spread/variation of heights as the range is larger 57 > 55 (with outlier) oe		B1ft
			(1)
(f)	36 ,, $a < x$ where 43 ,, x ,, 45 or 54 ,, $2a$,, 80		M1
	36 ,, a ,, "43" and 54 ,, $2a$,, 80		A1ft
	36 ,, a ,, 40		A1 (3)
Notes			Total 13
(a)	B1	Cao	
(b)	B1	for Q_1 correct	
	B1	for Q_3 correct	
(c)	M1	for either method correct or a correct value (ft their Q_1 and their Q_3)	
	A1	Both limits for outliers correct and statement about the outlier or the outlier given.	
(d)	M1	for a box drawn with only 2 whiskers, only one at each end (condone median line missing)	
	M1	for upper whisker ending at 76 (or 84 ft their upper outlier limit) and lower whisker ending at 28	
	M1	for Q_1 , Q_2 and Q_3 plotted, with $Q_2 = 48$ and ft their Q_1 and Q_3	
	A1	for a fully correct box plot with the outlier correctly shown – must be only 1	
(e)	B1ft	for a correct comment, referring to heights , with reference to a correctly named statistic . Must include the figures compared. Allow 'grow more/bigger' to imply taller Ignore any reference to skew SC If Q1/Q3 are incorrect then allow a ft comment about spread referring to the difference in IQR if compared to 18	
(f)	M1	for either range correct. Allow $72 \leq 2a \leq 80$ or $27 \leq a \leq 40$ for 54 ,, $2a$,, 80 Condone < rather than ,,	
		May be seen as separate inequalities e.g. $2a \leq 80$, $2a \geq 54$ is allowed for 54 ,, $2a$,, 80 A final answer of 36 ,, a ,, 40 or $36 < a < 40$ implies M1	
	A1ft	for both ranges correct ft their k . Allow $72 \leq 2a \leq 80$ or $27 \leq a \leq 40$ for 54 ,, $2a$,, 80 Condone < rather than ,, May be seen as separate inequalities	
	A1	Allow 36 to 40 or 36, 37, 38, 39, 40 NB It is possible to get M1A0A1	

Question	Scheme		Marks
2 (a)	$[P(2X - 3 > 5) =]0.45$		B1
			(1)
(b)	$E(X^2) = 2^2 \times 0.25 + 4^2 \times 0.3 + 5^2 \times 0.2 + 7^2 \times 0.1 + 8^2 \times 0.15 [= 25.3]$		M1
	$\text{Var}(X) = 2^2 \times 0.25 + 4^2 \times 0.3 + 5^2 \times 0.2 + 7^2 \times 0.1 + 8^2 \times 0.15 - 4.6^2$ or $\text{Var}(X) = "25.3" - 4.6^2$		M1
	$= 25.3 - 4.6^2 = 4.14 *$		A1*
			(3)
(c)	$[E(Y) =]13.4 = a \times 4.6 - b$		M1
	$[Var(Y) =]a^2 \times 4.14 = 66.24$		M1
	$a = 4$		A1
	$b = 5$		A1
			(4)
(d)	Sam throws 8 and Alex throws 2, 4 or 5 Sam throws 7 and Alex throws 2 or 4 Sam throws 5 or 4 or 2 and Alex throws 2		M1
	$0.15 \times (0.25 + 0.3 + 0.2)$ or $0.1 \times (0.25 + 0.3)$ or $(0.2 + 0.3 + 0.25) \times 0.25$		M1
	$0.15 \times (0.25 + 0.3 + 0.2) + 0.1 \times (0.25 + 0.3) + (0.2 + 0.3 + 0.25) \times 0.25$		M1
	$= 0.355 \left(= \frac{71}{200} \right)$		A1
			(4)
ALT	Alex throws 2 Alex throws 4 and Sam throws 7 or 8 Alex throws 5 and Sam throws 8		M1
	0.25 or $0.3 \times (0.1 + 0.15)$ or 0.2×0.15		M1
	$0.25 + 0.3 \times (0.1 + 0.15) + 0.2 \times 0.15$		M1
	$= 0.355 \left(= \frac{71}{200} \right)$		A1
Notes			Total 12
(a)	B1	0.45 oe	
(b)	M1	for a correct method to find $E(X^2)$ At least 3 terms correct and added This is not implied by 25.3 on its own	
	M1	for use of correct equation ft their $E(X^2)$	
	A1*	for a correct expression, with all terms seen, leading to the given answer	
(c)	M1	for writing or using a correct equation for $E(Y)$	
	M1	for writing or using a correct method for $\text{Var}(Y)$	
	A1	for $a = 4$ may be seen as part of the expression $4X \pm \dots$	
	A1	for $b = 5$ may be seen as part of the expression $\dots X - 5$	
(d)	M1	for recognising all the required combinations – implied by 3 correct probabilities Ignore any repeats but do not ignore any incorrect combinations	
	M1	for any one correct calculation from the 3 given	
	M1	for any 2 correct calculations from the 3 given	
	A1	0.355 oe	

Question	Scheme		Marks
3(a)	Width = 1.25 [cm]		B1
	18.75 cm ² for freq of 20 so $\frac{18.75}{20} \times 16 = 15$ cm ² for a frequency of 16 or $w \times h = 15$ or $fd = 5$		M1
	[$h = 15 \div 1.25$ or $h = 8 \div 5 \times 7.5 = 12$ (cm)]		A1
			(3)
(b)	$Q_2 = [32 +] \frac{7}{20} \times 4$ or using $n + 1$ gives $Q_2 = [32 +] \frac{7.5}{20} \times 4$		M1
	= 33.4 ($n + 1$ gives 33.5)		A1
			(2)
(c)	$\bar{y} = \frac{104}{50} [= 2.08]$	$\sum(w - 20) = 10 \times 104 [= 1040]$ or $\sum w = 10 \times 104 + 50 \times 20 [= 2040]$	M1
	$\bar{w} = 10 \times "2.08" + 20 = 40.8^*$	$\frac{"1040"}{50} + 20 = 40.8$ or $\frac{"2040"}{50} = 40.8$	A1*
			(2)
(d)	[Variance of $y =] \frac{233.54}{50} - ("2.08")^2 [= \frac{861}{2500} = 0.3444]$ or $10 \times \text{sd of } y = \text{sd of } w$		M1
	or $100 \times 233.54 = \sum(w^2) - 40 \times "2040" + 50 \times 400 [\Rightarrow \sum(w^2) = 84954]$ oe		
	[Variance of $w =] "0.3444" \times 100$ or $\frac{"84954"}{50} - 40.8^2 [= \frac{861}{25} = 34.44]$		M1
	or sd of $y = \sqrt{"0.3444"} [= \frac{\sqrt{861}}{50} = 0.5868\dots]$		
	sd of $w = \sqrt{"0.3444" \times 100}$ or $\sqrt{"34.44"}$ or $10 \times \frac{\sqrt{861}}{50}$	M1	
	= 5.868...	awrt 5.87	A1
(e)(i)	The mean would not change (as 40.8 is the mean)		B1
(ii)	The standard deviation would decrease (as 40.8 is in the middle so data closer together)		B1
	Both correct with a correct reason for why the standard deviation decreases		ddB1
			(3)
Notes			Total 14
(a)	B1	for width = 1.25 no need for units	
	M1	for sight of 15 or "their w " \times "their h " = 15 or $fd = 5$ May be implied by $h = 12$	
	A1	for height = 12 no need for units	
(b)	M1	for $\frac{7}{20} \times 4$ or $\frac{13}{20} \times 4$ or $\frac{m-32}{25-18} = \frac{4}{20}$ oe or $\frac{36-m}{38-25} = \frac{4}{20}$ oe (allow 25.5 rather than 25)	
	A1	33.4 or if using $(n + 1)$ 33.5	
(c)	M1	for a correct method to find the mean of y or $\sum(w - 20)$ or $\sum w$ ($10 \times 104 + k$ where $k \neq 20 \times 50$ is M0)	
	A1*	for a correct method to find mean of w which leads to 40.8	
(d)	M1	for a correct method to find the Variance of y or writing/using $10 \times \text{sd of } y = \text{sd of } w$ or a correct equation to find $\sum w^2$	
	M1	for a correct method to find the Variance of w or sd of y ft their $\text{Var}(y)$	
	M1	for a correct method to find the sd of w ft their $\text{Var}(w)$	
	A1	awrt 5.87 NB an exact answer $\frac{\sqrt{861}}{5}$ scores A0	
(e)(i)	B1	for no change (No reason needed) Allow mean = 40.8 to imply no change	
(ii)	B1	for sd decreases/be smaller/go down (condone Var decreases) (No reason needed)	
	ddB1	Both previous B1 awarded. For a correct reason for the sd decreasing Allow $(x - \bar{x}) = 0$ Allow $\sum(x - \bar{x})^2$ doesn't change and n increases. Allow the values would be more concentrated about the mean	

Question	Scheme		Marks
4 (a)	$S_{dg} = 141978.84 - \frac{1456.8 \times 713.2}{8}$ or $S_{gg} = 72675.98 - \frac{713.2^2}{8}$		M1
	$S_{dg} = 12105.12$		A1
	$S_{gg} = 9094.2$		A1
			(3)
(b)	$r = \frac{"12105.12"}{\sqrt{16769.78 \times "9094.2"}}$		M1
	$= 0.9802\dots$ awrt 0.98		A1
			(2)
(c)	$b = \frac{"12105.12"}{16769.78} [= 0.7218\dots]$		M1
	$a = \frac{713.2}{8} - "0.7218\dots" \times \frac{1456.8}{8} [= -42.297\dots]$		M1
	$g = -42.3 + 0.722d *$		A1*cso
			(3)
(d)	for each 1 [cm] increase in length/d the girth/g increases by "0.722..."		B1
			(1)
(e)(i)	138.2 awrt 138		B1
	(ii) [unreliable] as get a negative girth		B1
			(2)
(f)	$0.722x = 17.3$		M1
	$x = 23.96\dots$ awrt 24		A1
			(2)
Notes			Total 13
(a)	M1	for a correct expression for S_{dg} or S_{gg}	
	A1	for 12105.12 Allow $\frac{302628}{25}$	
	A1	for 9094.2 Allow $\frac{45471}{5}$	
		If exact answers are not seen then SC award M1A0A1 for both awrt 12100 and awrt 9090 if correct methods are seen If exact answers are not seen then SC award M1A0A0 for both awrt 12100 and awrt 9090 if no working seen	
(b)	M1	for a valid attempt at r with their S_{dg} not equal to 141978.84	
	A1	awrt 0.98	
(c)	M1	for a correct method to find the value of b May be implied by 0.7218 or better	
	M1	for a correct method to find a ft their b May be implied by -42.29 or better	
	A1*	both method marks must be awarded with sight of 0.7218 or better or -42.29 or better	
(d)	B1	for a suitable contextual comment that mentions 0.722 (or better) If units are stated they must be correct.	
(e)(i)	B1	awrt 138 Allow 1.38m	
(ii)	B1	For a correct reason eg sd = 45.8 cm so girth is nearly 3sd below mean so likely outlier Allow substitution of 50 leading to $g = -6.2$ and suitable reason e.g. this is not possible/this is negative Do not allow substitution of 0.5 to imply the girth is negative	
(f)	M1	for a correct equation. implied by awrt 24	
	A1	awrt 24	

Question	Scheme		Marks
5 (a)	$P(X < 18) = P\left(Z < \pm\left(\frac{18-15}{2}\right)\right) [= \pm 1.5]$		M1
	= 0.9332 awrt 0.933		A1
			(2)
(b)	$\frac{x-15}{2} = 0.2533$		M1B1
	$x = 15.506\dots$ awrt 15.5		A1
			(3)
(c)	$P(T > \mu - 10) = 0.975$		M1
	$\frac{(\mu \pm 10) - \mu}{\sigma} = \pm 1.96 \Rightarrow \sigma = \frac{10}{1.96} [= 5.10\dots]$		M1
	$P(T > \mu - 5) = P\left(Z > \frac{\mu - 5 - \mu}{"5.10\dots"} [= -0.98]\right) [= 0.836\dots]$		M1
	$P(T > \mu - 5 \mid T > \mu - 10) = \frac{"0.836\dots"}{"0.975"}$		M1
	= 0.8579... awrt 0.858		A1
			(5)
Notes			Total 10
(a)	M1	for standardising correctly May be implied by ± 1.5	
	A1	awrt 0.933 (Do not ISW)	
(b)	M1	for correct standardisation = to a z value such that 0.25 ,, $ z $,, 0.26	
	B1	for use of awrt ± 0.2533	
	A1	awrt 15.5	
(c)	M1	for the correct probability of 0.975 – may be seen as the denominator of the fraction. May be implied by use of $ z = 1.96$ or better	
	M1	For $\frac{\mu + 10 - \mu}{\sigma} = 1.96$ or $\frac{\mu - 10 - \mu}{\sigma} = -1.96$, leading to a value for σ May be implied by ± 0.98	
	M1	for a correct method to find $P(T > \mu - 5)$ using their value for σ May be implied by -0.98 If $P(T < \mu + 5)$ is calculated then this may be implied by 0.98	
	M1	for $\frac{p}{0.975}$ where $0.5 < p < 0.975$ (must be a probability not their z value) If the denominator is incorrect only follow through their $P(T > \mu - 10)$ if clearly labelled and > 0.95	
	A1	awrt 0.858	

Question	Scheme		Marks
6(a)	0.16 oe		B1 (1)
(b)	Mark parts (b) and (c) together If values are given in the diagram and the script, then the script takes precedence		
	$[P(C)=]0.04+0.15+0.12+p[=0.31+p]$	$[P(C')=]0.1+0.23+q[=0.33+q]$	M1M1
	$[P(S)=]0.1+0.15+0.12+0.23[=0.6]$	$[P(S')=]1-(0.12+0.15+0.1+0.23)[=0.4]$	
	$[P(S \cap C)=]0.15+0.12[=0.27]$	$[P(S' \cap C')=]q$	M1d
	$("0.31"+p) \times "0.6" = "0.27" \text{ oe}$	$("0.33"+q) \times "0.4" = q \text{ oe}$	A1
	$p = 0.14 \text{ oe}$	$q = 0.22 \text{ oe}$	(4)
(c)	$q = 1 - (0.04 + 0.12 + 0.15 + 0.1 + 0.23 + "p")$	$p = 1 - (0.04 + 0.12 + 0.15 + 0.1 + 0.23 + "q")$	M1
	$q = 0.22 \text{ oe}$	$p = 0.14 \text{ oe}$	A1ft (2)
(d)(i)	$[P((C \cup S) \cap G')] = 0.39 \text{ oe}$		B1 (1)
(ii)	$P(C (S \cap G)) = \frac{0.15}{0.15+0.1}$		M1
	$= 0.6 \text{ oe}$		A1 (2)
(e)	Number of teenagers = $\frac{76}{0.15+"p"} \text{ oe}$		M1
	Number who don't play Scrabble = $"\left(\frac{76}{0.15+p}\right)" \times 0.4 \text{ (=104.8...)}$		M1
	$= 104.8... \text{ awrt } 105$		A1 (3)
Notes			Total 13
(a)	B1	correct probability	
(b)	M1	for 2 correct probability expressions	
	M1	all 3 correct probability expressions Allow $P(C) = 0.45$	
	M1d	dependent on the 1st M1 being awarded for use of $P(C \cap S) = P(C) \times P(S)$ oe or $P(C' \cap S') = P(C') \times P(S')$ ft their probabilities if labelled clearly	
	A1	for 0.14 or exact equivalent or 0.22 or exact equivalent	
(c)	M1	for a correct expression for q ft their value of p or a correct expression for p ft their value of q May be implied by a correct value for q ft their p or a correct value for p ft their q	
	A1ft	for 0.22 or exact equivalent ft their value of p or 0.14 or exact equivalent ft their value of q ($p + q = 0.36$ provided p and q are probabilities)	
(d)(i)	B1	for 0.39 or exact equivalent do not allow $0.04 + 0.12 + 0.23$	
(ii)	M1	for $\frac{0.15}{0.15+0.1}$	
	A1	for 0.6 or exact equivalent	
(e)	M1	Relating 76 to their $P(C \cap G)$ May be implied by awrt 262	
	M1	for number of teenagers $\times 0.4$ ft their number of teenagers e.g. $0.4 \times "262"$ provided that the number of teenagers is not 76	
	A1	awrt 105 ISW	

