

Mark Scheme (Results)

January 2024

Pearson Edexcel International Advanced Subsidiary Level in Chemistry (WCH13) Paper 01 Practical Skills in Chemistry I

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <u>www.edexcel.com</u> or <u>www.btec.co.uk</u>. Alternatively, you can get in touch with us using the details on our contact us page at <u>www.edexcel.com/contactus</u>.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2024 Question Paper Log Number P75593A Publications Code WCH13_01_2401_MS All the material in this publication is copyright © Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

() means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the <u>meaning</u> of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|---|---|------|
| 1(a) | An answer that makes reference to the following points: | Allow alkanones | (3) |
| | (Functional group 1) ketone (1) | Ignore carbonyl compound Do not award aldehyde | |
| | (Functional group 2) alkene (1) | Ignore carbon – carbon double bond Do not award alkyl | |
| | (Functional group 3) (secondary) alcohol (1) | Allow hydroxy / hydroxyl Do not award hydroxide Do not award primary / tertiary | |
| | | Ignore cyclo / aliphatic / aromatic in M1 to M3 Ignore any formulae Allow names shown on the diagram for M1 to M3 | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 1(b) | An answer that makes reference to the following points: • Functional group 2 / alkene Test – (add) bromine water / Br ₂ (aq) (1) Result – (turns from orange to) colourless / (yellow to) colourless (1) | M2 dependent on M1 or near miss – e.g. omission of 'acidified' for dichromate(VI) test means M1 cannot be awarded, but M2 is still available. Allow bromine / Br ₂ Allow (brown to) colourless / (brown-red to) colourless Allow just decolourises Do not award incorrect initial colour e.g. red to colourless Allow Test - add acidified KMnO ₄ / (potassium) manganate ((VII)) (1) Result – (turns from purple/pink to) colourless/decolourises (1) | (4) |
| | Functional Group 3 / 2° alcohol EITHER Test – (add) PCl₅ / phosphorus pentachloride / phosphorus(V) pentachloride (1) Result – steamy / misty fumes (1) OR Test – (add) Na (1) Result - Bubbles (of colourless gas) (1) | Allow (add) SOCl ₂ /PCl ₃ Allow white fumes / fumes that turn indicator paper red / fumes that form white smoke with NH ₃ Allow gas / vapour as alternative to fumes Ignore references to HCl Ignore white smoke without reference to NH ₃ Allow Test – (add) acidified dichromate ((VI)) Result - orange to green Allow Test (add) carboxylic acid and H ⁺ and warm/heat Result - fruity smell | |
| | Comment – ignore reference to products in Na test even if incorrect | | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---------------------|------|
| 1(c)(i) | An answer that makes reference to the following point: | Ignore any formulae | (1) |
| | • 96 / ninety-six | | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 1(c)(ii) | An answer that makes reference to the following point: • (free) radical | Allow R' Allow 'particle with an unpaired electron' | (1) |

(Total for Question 1 = 9 marks)

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|---|---|------|
| 2(a) | An answer that makes reference to the following points: both '24.30' and '24.20' circled | Allow other forms of indication e.g. use of asterisks / underlining / circling of '1' and '3' in titration header Do not award any other values | (1) |

| Titration | Rough | 1 | 2 | 3 |
|---|-------|-------|-------|-------|
| Burette reading (final) / cm ³ | 24.60 | 48.90 | 23.80 | 48.00 |
| Burette reading (initial) / cm ³ | 0.00 | 24.60 | 0.00 | 23.80 |
| Titre / cm ³ | 24.60 | 24.30 | 23.80 | 24.20 |

| Question | Acceptable Answers | Additional Guidance | Mark |
|----------|---|--|------|
| Number | | | |
| 2(b) | | Example of calculation | (1) |
| | • calculation of mean from titres 1 and | - | |
| | 3 to 4 SF (even if not circled) | $(24.30 + 24.20) \div 2 =$ | |
| | | 24.25 (cm ³) Accept 0.02425 dm ³ | |
| | | Allow TE from (a) provided they are calculating the mean of more than one circled titre | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|-----------------------------------|------|
| 2(c)(i) | An answer that makes reference to the following point:colourless to (pale) pink | Do not award colourless to purple | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|---|--|------|
| 2(c)(ii) | calculation of the amount of NaOH (in mol) (1) calculation of moles of citric acid in 25 cm³ of diluted lemon juice (1) calculation of moles of citric acid in 250 cm³ of diluted lemon juice (1) calculation of mass of citric acid (1) calculation of percentage by mass of citric acid in lemon juice and to 2 SF (1) | Example of calculation Ignore rounding errors in M1 to M4 Ignore SF in M1 to M4 unless 1SF $(24.25 \div 1000) \times 0.103 = 2.4978 \times 10^{-3}/0.0024978 \text{ (mol)}$ Allow TE from (b) $2.4978 \times 10^{-4} \div 3 = 8.3258 \times 10^{-4}$ /0.0083258 (mol) $8.3258 \times 10^{-4} \times 10 = 8.3258 \times 10^{-3}$ /0.0083258 (mol) $8.3258 \times 10^{-3} \times 192 = 1.5986 \text{ (g)}$ M1, M2, M3 and M4 can be in any order $((1.5986 \div 24) \times 100 = 6.6608 \% =)6.7 (\%)$ Correct answer scores 5 marks Allow TE from M4 if answer is less than 100% Ignore units in intermediate stages even if incorrect | (5) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 2(d) | An answer that makes reference to the following point: the other acids are present in (very) small amounts (in lemon juice, so will have little effect on the titre values) | Allow 'citric acid has a much greater concentration' Allow 'the other acids are present in much smaller amounts' Ignore just 'the other acids are present in smaller amounts' Ignore comments related to pH / acid strength | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 2(e)(i) | An answer that makes reference to the following point: (pour away solution, rinse flask and) make a new / fresh solution (of diluted lemon juice) | Allow start again (from the beginning) / repeat the procedure / do the experiment again Allow transfer to larger (volumetric) flask, (rinse) and make up to volume | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|-------------|
| 2(e)(ii) | An answer that makes reference to the following points: because some water has evaporated (1) (mean NaOH) titre would be lower (as NaOH (aq) is more concentrated) (1) OR NaOH (reacts with CO₂ to) form Na₂CO₃ / NaHCO₃ (1) which also reacts with the acid, so will not affect the titre / which results in a less sharp end point (1) OR because NaOH reacts with CO₂ (in the air) (1) (mean) titre would be greater (as NaOH (aq) is less concentrated) (1) | M2 dependent on M1 Allow water has escaped / been lost Allow some of the solution may have evaporated Do not award some of the NaOH (solution) has evaporated Comment If no other mark is awarded allow 'the bottle is sealed so the titre would not be different' for 1 mark | (2) |
| | | $\frac{ }{(Total for Ouestion 2 = 1)}$ | 2 marks |

(Total for Question 2 = 12 marks)

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|------|
| 3(a) | An answer that makes reference to the following point: | | (1) |
| | • the compounds containing iodine are ionically bonded | Allow the compounds containing iodine do not contain hydrogen / carbon Allow iodine present as iodide ions Ignore iodine compounds are | |
| | Comment 'It' or 'they' can be assumed to refer to the compounds containing iodine | inorganic / not organic Allow the compounds containing iodine have high(er) boiling temperatures (so do not vaporise readily) Ignore references to bond energy / solubility in water / intermolecular forces / melting point | |
| | | Do not award iodine has a high boiling temperature | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|------|
| 3(b) | calculation of moles of oxygen produced by 1 dm³ of hydrogen peroxide solution (1) calculation of concentration of hydrogen peroxide (1) | Example of calculation $20 \div 24 = 0.83333/8.3333 \times 10^{-1}$ (mol) $(0.83333 \times 2) = 1.6667 = 1.67$ $(mol dm^{-3})$ Allow TE from M1 to M2Ignore SF except 1 SFIgnore minor slips in units e.g.mol dm ³ or mol/dm ⁻³ Do not award major unit errors inM2 e.g. g or dm ³ mol ⁻¹ | (2) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 3(c)(i) | An answer that makes reference to the following points: | Allow multiples | (2) |
| | • (Oxidation of iodide ions) $2I^- \rightarrow I_2 + 2e^{(-)}$ (1) | Accept $2I^ 2e^{(-)} \rightarrow I_2$ | |
| | (Reduction of hydrogen peroxide under acidic conditions) H₂O₂ + 2H⁺ + 2e⁽⁻⁾ → 2H₂O (1) | Allow 2H ⁺ above arrow Ignore state symbols even if incorrect | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|------|
| 3(c)(ii) | An answer that makes reference to the following point: | Allow multiples | (1) |
| | • $H_2O_2 + 2H^+ + 2I^- \rightarrow 2H_2O + I_2$ | Allow 2H ⁺ above arrow Allow 2HI Ignore state symbols even if incorrect | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|------|
| 3(c)(iii) | An answer that makes reference to the following point: • yellow / brown (solution) | Ignore adjectives e.g. 'pale' Do not award orange / purple / blue / black / red / pink / red-brown Do not award precipitate / ppt / solid / crystals Do not award vapour / gas / fumes | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|---|---|------|
| 3(d) | An answer that makes reference to the following points: add (aqueous) filtrate / iodine (solution) / mixture to (separating funnel containing) cyclohexane (1) (gently) shake / invert separating funnel (1) open tap of (inverted) separating funnel occasionally to relieve pressure / release gas (1) (allow layers to settle) and then remove (lower) aqueous / inorganic layer (1) | Allow addition of liquids to separating funnel in any order Ignore minor slips with name of solvent e.g. cyclohexene / hexane / cyclohexanol Allow idea of mixing / swirling contents Ignore stirring Accept loosen/remove stopper (slightly) to relieve pressure / release gas Ignore references to drying agents / Na ₂ CO ₃ / NaHCO ₃ Allow removal of the lower layer if it is clear that it is not the cyclohexane layer Allow run off the water layer Ignore any references to colours Ignore subsequent attempts to remove iodine from cyclohexane Do not award removal of cyclohexane before aqueous layer M1 to M4 could be shown on an annotated diagram | (4) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 3(e)(i) | An answer that makes reference to the following point: • (serious) health hazard | Accept hazard to health Allow can cause serious health damage Ignore risk to health / specific conditions e.g. damage to lungs / causes respiratory problems Do not award toxic / | (1) |
| | | corrosive / explosive | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|---|---|------|
| 3(e)(ii) | An answer that makes reference to the following points: | | (2) |
| | carry out in fume cupboard (1) | Allow carry out in a well- ventilated room Ignore wear goggles, gloves, lab coats and masks Ignore use small amount (of cyclohexane) | |
| | avoid contact with (naked) flames / sources of ignition (1) | Allow don't put near fire / don't use near a Bunsen (burner) / don't put near flame Ignore keep away from heat / use of an electric heater / water bath | |

(Total for Question 3 = 14 marks)

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 4(a)(i) | • calculation of density of ethanol- water mixture sample A | Example of calculation $4.75 \div 5.00 = 0.95(0) \text{ (g cm}^{-3})$ Ignore units even if incorrect Allow 950 g dm ⁻³ | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 4(a)(ii) | An answer that makes reference to the following points: • suitable linear scale with points covering at least half the available space in both directions (1) • all points plotted accurately within $\pm \frac{1}{2}$ a square (1) • suitable curve of best fit (1) | $\begin{array}{c} 0.98\\ 0.96\\ 0.94\\ 0.92\\ 0.9\\ 0.9\\ 0.88\\ 0.86\\ 0.84\\ 0.82\\ 0.8\\ 30 & 50 & 70 & 90 & 110 \end{array}$ | (3) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 4(a)(iii) | An answer that makes reference to the following point: answer from (a)(i) used to determine percentage with evidence shown on graph Comment evidence could be horizontal and / or vertical line OR additional point plotted at density from (a)(i) | 38.5 (%) Answer must be consistent with their line on graph ± ½ a square Allow TE from straight line graph Ignore units even if incorrect | (1) |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|--|------|
| 4(b)(i) | An explanation that makes reference to the following points: (distillation allows) removal of ethanal / aldehyde /product (as it forms) (1) to prevent formation of a carboxylic acid / ethanoic acid (1) | Allow ethanal can be distilled off / distilled out Allow aldehyde is separated Ignore just distillation Allow heating under reflux forms a carboxylic acid Allow to prevent further oxidation Allow it is only oxidised to an aldehyde Do not award to prevent formation of / further | (2) |
| | | oxidation to a ketone | |

| Question Number | Acceptable Answers | Additional Guidance | Mark |
|--------------------|--|---|------|
| 4(b)(ii) | An explanation that makes reference to the following points: prevents (formation of) air gaps / bubbles / air spaces (1) (so more) efficient cooling results (1) | Allow so water fills it / condenser Allow maximum cooling / faster cooling / better cooling Allow as bubbles reduce the cooling effect Ignore references to condensing | (2) |

| Question Number | Acceptable Answers | Addi | tional Guidance | Mark |
|--------------------|--|--|---|------|
| 4(c)(i) | • calculation of energy transferred to water | | Example of calculation $(100 \times 4.18 \times (57.9 - 20.0)) =$ | |
| | | | 2.2 (J) w 15.842 kJ | |
| | | - | re SF except 1 SF re any positive or negative | |
| Question Number | Acceptable Answers | 0 | tional Guidance | Mark |
| 4(c)(ii) | | Exan | ple of calculation | (1) |
| | • calculation of amount of ethanol burnt in mol | | $(0.650 \div 46.0 =)$ 0.014130 / 1.4130 × 10 ⁻² / 14.130 × 10 ⁻³ (mol) | |
| | | Ignore SF except 1SF | | |
| Question Number | Acceptable Answers | Additional Guidance | | Mark |
| 4(c)(iii) | | Example of calculation | | (1) |
| | calculate the enthalpy change of combustion of ethanol in kJ mol⁻¹ | (15.8 | (15.8422 ÷ 0.014130 = (-) 1121.2) | |
| | and | $= -1120/1100 (kJ mol^{-1})$ | | |
| | negative sign and | Allow TE from (c)(i) and (c)(ii) Allow answer consistent with | | |
| | 3SF or 2SF | rounded value from (c)(ii) e.g. use | | |
| | | of 0.014 gives $-1131.585 = -1130 (kJ mol^{-1})$ | | |
| | | Allow -1120000 / -1100000 J mol ⁻¹ | | |
| Question Number | Acceptable Answers | | Additional Guidance | Mark |
| 4(d) | An explanation that makes reference to the following points: | | Mark independently | (3) |
| | as copper coil conducts heat from emissions / waste gases / hot gases into water (1) | | Allow lid prevents / reduces evaporation of water Ignore comments about the stirrer / surface area of copper tube | |

| less heat loss (to the atmosphere) (1) (constant) supply of air into apparatus so less chance of incomplete combustion (1) | Allow 'no heat loss' Allow 'less energy lost' Allow oxygen / air prevents incomplete combustion' Allow oxygen / air allows complete combustion to occur |
|---|---|
| | Allow reverse argument if it is clear the copper can is being referred to |

(Total for Question 4 = 15 marks)

FIRST RELEASED ON AP - EDEXCEL DISCORD https://sites.google.com/view/ap-edexcel

TOTAL FOR PAPER = 50 MARKS

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom