

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	

Pearson Edexcel International Advanced Level

Thursday 23 January 2025

Morning (Time: 1 hour 20 minutes)

Paper reference **WCH13/01**

Chemistry

International Advanced Subsidiary/Advanced Level

UNIT 3: Practical Skills in Chemistry I

You must have:
Scientific calculator, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P78457A

©2025 Pearson Education Ltd.
H:1/1/1/1/




Pearson

Answer ALL the questions. Write your answers in the spaces provided.

1 This question is about qualitative testing for ions.

(a) A student was given a solid sample of potassium sulfate, K_2SO_4 .

- (i) Describe how the student could use flame colour to confirm that the solid sample contains potassium ions.

Outline the experimental procedure, stating the colour of the flame produced.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) The student dissolved some of the solid sample in deionised water in a test tube.

To the test tube, the student added dilute hydrochloric acid followed by aqueous barium chloride. A precipitate was observed.

Give the colour of the precipitate and the **ionic** equation for its formation. Include state symbols.

(2)

Colour of precipitate

.....

Ionic equation



- (b) Another student was given a solution containing one type of cation and two types of anion.
- (i) The student warmed some of the solution in a test tube, before adding aqueous sodium hydroxide and holding damp red litmus paper at the mouth of the tube.

From the results, the student identified the cation as the ammonium ion, NH_4^+ .

Explain the student's inference.

(2)

- (ii) To another test tube of the solution, the student added silver nitrate solution. A pale yellow precipitate was observed.

The student then added an **excess** of dilute nitric acid. Effervescence was observed.

Some of the precipitate dissolved.

The student added an excess of **concentrated** ammonia solution to the remaining precipitate.

The precipitate dissolved to form a colourless solution.

Complete the table about the student's results.


(3)

Formula of gas evolved	
Formula of the precipitate that dissolved in dilute nitric acid	
Formula of the precipitate that dissolved in concentrated ammonia solution	

(Total for Question 1 = 11 marks)



- 2 A student used the reaction of magnesium with hydrochloric acid to determine the volume of 1 mol of hydrogen gas at room temperature and pressure (r.t.p.).



Procedure

Step 1 Weigh 0.030 g of magnesium ribbon.


Step 2 Add approximately 25 cm³ of 2.0 mol dm⁻³ hydrochloric acid into a burette and carefully add 25 cm³ of water on top.

Step 3 Push the magnesium ribbon into the open end of the burette so that it stays in position.

Step 4 Quickly invert the burette. Clamp the burette vertically and take the burette reading.

Step 5 Wait for the magnesium to start reacting. Take the final burette reading after all the magnesium has reacted.

A diagram of the experiment is shown.



- (a) Explain why the volume of hydrochloric acid added to the burette in Step **2** does not need to be exact.

You must include a calculation.

(3)

.....

.....

.....

- (b) State why the magnesium does not start reacting as soon as the burette is inverted in Step **4**.

(1)

.....


.....

.....

.....



(c) The diagram shows the burette readings from Steps 4 and 5.




Record the burette readings, and hence the volume of hydrogen gas produced in the experiment.

(2)

Burette reading in Step 4 / cm ³	
Burette reading in Step 5 / cm ³	
Volume of hydrogen gas / cm ³	

(d) Calculate the volume of 1 mol of hydrogen gas at r.t.p., in cm³, using your answer to (c), the reaction equation and the mass of magnesium (0.030 g) used in the procedure.

(2)



- (e) The magnesium ribbon used in Step 1 contained a magnesium oxide impurity of up to 2% by mass.

Explain whether this magnesium oxide impurity could account for any difference in your answer to (d) from the Data Book value of $24\,000\text{ cm}^3$.

No calculation is required.

(2)

- (f) Two readings were taken to measure the mass of magnesium used in the experiment. The uncertainty in each reading was 0.0025 g .

- (i) Calculate the percentage uncertainty in the mass of magnesium used in the experiment.

(1)

- (ii) The student wanted to reduce the percentage uncertainty by using double the mass of magnesium.

Give the reason why the experiment outlined in the procedure could **not** be conducted successfully using double the mass of magnesium.

(1)

(Total for Question 2 = 12 marks)



3 The concentration of a solution of barium hydroxide can be determined using a standard solution of ethanedioic acid.

(a) State what is meant by the term **standard** solution.

(1)

(b) Describe the preparation of 100.0 cm^3 of a $0.0500\text{ mol dm}^{-3}$ standard solution of ethanedioic acid, starting with crystals of ethanedioic acid dihydrate, $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$.

Your answer should include details of the apparatus used and the mass of $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ required.

(6)



(c) 25.0 cm^3 of $0.0500\text{ mol dm}^{-3}$ ethanedioic acid solution, $\text{H}_2\text{C}_2\text{O}_4(\text{aq})$, was added to a conical flask and titrated with barium hydroxide solution, $\text{Ba}(\text{OH})_2(\text{aq})$, of an unknown concentration.

- (i) Name the piece of apparatus that should be used to add the ethanedioic acid solution, $\text{H}_2\text{C}_2\text{O}_4(\text{aq})$, to the conical flask.

(1)

- (ii) The titration was carried out with a few drops of phenolphthalein indicator in the conical flask.

State the colour change observed at the end-point of the titration.

(2)

From to

- (iii) The formation of a white precipitate was observed during the titration.

Suggest the identity of the white precipitate, by name or formula.

(1)

- (iv) The titration was repeated, giving a mean titre of 31.55 cm^3 .

Calculate the concentration of the barium hydroxide solution, $\text{Ba}(\text{OH})_2(\text{aq})$, in g dm^{-3} .

$\text{Ba}(\text{OH})_2$ and $\text{H}_2\text{C}_2\text{O}_4$ react in a 1:1 ratio.

Give your answer to an appropriate number of significant figures.


(3)

(Total for Question 3 = 14 marks)



P 7 8 4 5 7 A 0 9 1 6

- 4 This question is about the preparation of 2-chloro-2-methylpropane from 2-methylpropan-2-ol and concentrated hydrochloric acid.



Procedure

- Step 1 Pour 9.0 cm^3 of 2-methylpropan-2-ol into a separating funnel.
- Step 2 Add 20 cm^3 of concentrated hydrochloric acid to the separating funnel, about 3 cm^3 at a time.
Stopper and invert the funnel after each addition.
- Step 3 Leave the separating funnel for about 20 minutes, shaking it gently at regular intervals. Allow the layers to separate.
- Step 4 Discard the aqueous layer before washing the organic layer repeatedly with sodium hydrogencarbonate solution, releasing the pressure after each addition.
- Step 5 Transfer the organic layer into a small, dry conical flask and add three spatula measures of anhydrous sodium sulfate, swirling the flask after each addition.
- Step 6 Decant the organic liquid into a pear-shaped flask and distil the 2-chloro-2-methylpropane into a pre-weighed conical flask.
- (a) Give **one** reason, relating to safety, why the hydrochloric acid is added to the separating funnel 3 cm^3 at a time, and not all at once, in Step 2.

(1)

.....

.....

.....



(b) Draw a labelled diagram of the separating funnel at the end of Step 3.

[Density of 2-chloro-2-methylpropane = 0.85 g cm^{-3}]

(2)

(c) Sodium hydrogencarbonate solution is added to remove acid impurities in Step 4.

Give the reason why a stronger alkali, such as sodium hydroxide solution, should **not** be used for this purpose.

(1)


(d) State the role of the sodium sulfate in Step 5.

(1)



- (e) Identify **two** changes that must be made to the apparatus shown before distilling in Step 6, giving a reason for each change.

(4)



Change 1

.....

Reason 1

.....

.....

Change 2

.....

Reason 2

.....

.....

- (f) Calculate the mass of 2-chloro-2-methylpropane produced in this procedure, if the percentage yield is 70%.

[Density of 2-methylpropan-2-ol = 0.78 g cm^{-3}]

[M_r of 2-methylpropan-2-ol = 74.0]

[M_r of 2-chloro-2-methylpropane = 92.5]

(4)

(Total for Question 4 = 13 marks)

TOTAL FOR PAPER = 50 MARKS



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1	2	3	4	5	6	7	0 (8)
6.9 Li lithium 3	9.0 Be beryllium 4	10.8 B boron 5	12.0 C carbon 6	14.0 N nitrogen 7	16.0 O oxygen 8	19.0 F fluorine 9	20.2 Ne neon 10
23.0 Na sodium 11	24.3 Mg magnesium 12	27.0 Al aluminium 13	28.1 Si silicon 14	31.0 P phosphorus 15	32.1 S sulfur 16	35.5 Cl chlorine 17	39.9 Ar argon 18
39.1 K potassium 19	40.1 Ca calcium 20	65.4 Zn zinc 30	63.5 Cu copper 29	58.7 Ni nickel 28	58.9 Co cobalt 27	57.8 Fe iron 26	54.9 Mn manganese 25
85.5 Rb rubidium 37	87.6 Sr strontium 38	112.4 Cd cadmium 48	107.9 Ag silver 47	106.4 Pd palladium 46	102.9 Rh rhodium 45	101.1 Ru ruthenium 44	[98] Tc technetium 43
132.9 Cs caesium 55	137.3 Ba barium 56	200.6 Hg mercury 80	197.0 Au gold 79	195.1 Pt platinum 78	192.2 Ir iridium 77	190.2 Os osmium 76	186.2 Re rhenium 75
[223] Fr francium 87	[226] Ra radium 88	204.4 Tl thallium 81	204.4 Pb lead 82	207.2 Bi bismuth 83	209.0 Po polonium 84	210 At astatine 85	[222] Rn radon 86

1.0
H
hydrogen
1

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140 Ce cerium 58	141 Pr praseodymium 59	144 Nd neodymium 60	147 Pm promethium 61	150 Sm samarium 62	152 Eu europium 63	157 Gd gadolinium 64	159 Tb terbium 65	163 Dy dysprosium 66	165 Ho holmium 67	167 Er erbium 68	169 Tm thulium 69	173 Yb ytterbium 70	175 Lu lutetium 71
232 Th thorium 90	[231] Pa protactinium 91	238 U uranium 92	[237] Np neptunium 93	[242] Pu plutonium 94	[243] Am americium 95	[247] Cm curium 96	[245] Bk berkelium 97	[251] Cf californium 98	[254] Es einsteinium 99	[253] Fm fermium 100	[256] Md mendelevium 101	[254] No nobelium 102	[257] Lr lawrencium 103

* Lanthanide series

* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

