

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

## Pearson Edexcel International Advanced Level

**Thursday 29 May 2025**

Morning (Time: 1 hour 30 minutes)

Paper  
reference

**WMA13/01**



### Mathematics

#### International Advanced Level

#### Pure Mathematics P3

#### You must have:

Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations.  
Calculators must not have the facility for symbolic algebra manipulation,  
differentiation and integration, or have retrievable mathematical formulae  
stored in them.**

#### Instructions:

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need*.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

#### Information:

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question*.

#### Advice:

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

**Turn over** ►

P76390A

©2025 Pearson Education Ltd.  
Y:1/1/1/1/



**Pearson**

- 1: In this question you must show all stages of your working.  
Solutions relying entirely on calculator technology are not acceptable.

The functions  $f$  and  $g$  are defined by

$$f(x) = \frac{2x}{3x+1} \quad x \in \mathbb{R} \quad x \geq 0$$

$$g(x) = 4 - x^2 \quad x \in \mathbb{R} \quad x \geq 0$$

- (a) Find the value of  $gf(1)$  (2)
- (b) Find the range of  $f$  (2)
- (c) Find  $f^{-1}(x)$  (2)
- (d) Solve  $f^{-1}(x) = f(x)$  (2)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

### **Question 1 continued**



## **Question 1 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



**DO NOT WRITE IN THIS AREA**

**DONOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

### **Question 1 continued**

**(Total for Question 1 is 8 marks)**



**2:** In this question you must show all stages of your working.  
Solutions relying entirely on calculator technology are not acceptable.

$$f(x) = 7 \cos x - 24 \sin x$$

- (a) Express  $f(x)$  in the form  $R \cos(x + \alpha)$  where  $R$  and  $\alpha$  are constants,  $R > 0$

and  $0 < \alpha < \frac{\pi}{2}$

Give the exact value of  $R$  and give the value of  $\alpha$ , in radians, to 3 decimal places.

(3)

$$g(x) = \frac{5}{90 - 3f(2x)}$$

- (b) Using the answer to part (a), find

- (i) the minimum value of  $g(x)$ , giving your answer as a fully simplified fraction,
- (ii) the smallest positive value of  $x$  for which this minimum value occurs, giving your answer to 3 decimal places.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**DO NOT WRITE IN THIS AREA**

## **Question 2 continued**

**(Total for Question 2 is 7 marks)**



P 7 6 3 9 0 A 0 7 3 2

3:


**Figure 1**

Figure 1 shows a linear relationship between  $\log_{10} y$  and  $\log_{10} x$ .

The line passes through the points  $(-3.5, 0)$  and  $(0, -2)$  as shown.

- (a) Find an equation linking  $\log_{10} y$  with  $\log_{10} x$  (2)
- (b) Hence, or otherwise, express  $y$  in the form  $px^q$  where  $p$  and  $q$  are rational constants. (3)



**DO NOT WRITE IN THIS AREA**

**DONOT WRITE IN THIS AREA**

### **Question 3 continued**

**(Total for Question 3 is 5 marks)**



P 7 6 3 9 0 A 0 9 3 2

4: The function  $f$  is defined by

$$f(x) = \frac{49x}{x^2 + x - 12} + \frac{7x}{x + 4} \quad x > 3$$

(a) Show that

$$f(x) = \frac{7x}{x - 3} \quad x > 3 \quad (3)$$

(b) Hence find  $f'(x)$  giving your answer in simplest form. (2)



## **Question 4 continued**

**(Total for Question 4 is 5 marks)**



## 5: Find

$$(i) \int \sin^2 3x \, dx \quad (2)$$

$$(ii) \int x(x^2 + 4)^{\frac{3}{2}} dx \quad (2)$$



**DO NOT WRITE IN THIS AREA**

[View Details](#) | [Edit](#) | [Delete](#)



P 7 6 3 9 0 A 0 1 3 3 2

13

*Turn over* 

- 6: In this question you must show all stages of your working.  
Solutions relying entirely on calculator technology are not acceptable.

The temperature,  $\theta^{\circ}\text{C}$ , of a computer processor,  $t$  minutes after the computer is switched off, is modelled by the equation

$$\theta = 21 + Ae^{-kt}$$

where  $A$  and  $k$  are positive constants.

Given that the temperature of the processor was  $75^{\circ}\text{C}$  when the computer was switched off,

- (a) find the value of  $A$ .

(2)

Given also that it takes 5 minutes for the temperature of the processor to decrease from  $75^{\circ}\text{C}$  to  $25^{\circ}\text{C}$ ,

- (b) find the value of  $k$ , giving your answer to 3 significant figures.

(3)

At time  $T$  minutes, the temperature of the processor is decreasing at a rate of  $9^{\circ}\text{C}$  per minute.

- (c) Find the value of  $T$  according to the model, giving your answer to 2 decimal places.

(3)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

## **Question 6 continued**



## **Question 6 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



## **Question 6 continued**

(Total for Question 6 is 8 marks)



7: A continuous curve has equation

$$y = e^{-x^2} \sin 3x \quad 0 \leq x \leq \frac{\pi}{3}$$

The curve has a stationary point at the point  $P$ .

- (a) Show, using calculus, that the  $x$  coordinate of  $P$  is a solution of the equation

$$x = \frac{1}{3} \arctan\left(\frac{3}{2x}\right) \quad (4)$$

Using the iteration formula

$$x_{n+1} = \frac{1}{3} \arctan\left(\frac{3}{2x_n}\right) \quad x_1 = 0.4$$

- (b) find the value of

(i)  $x_2$

(ii)  $x_4$

giving your answers to 4 decimal places.

(3)

- (c) Using a suitable interval and a suitable function which should be stated, show that the  $x$  coordinate of  $P$  is 0.430 correct to 3 decimal places.

(2)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

## **Question 7 continued**



## **Question 7 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



## **Question 7 continued**

(Total for Question 7 is 9 marks)



**8:** In this question you must show all stages of your working.  
Solutions relying entirely on calculator technology are not acceptable.

(a) Prove that

$$\tan 3x \equiv \frac{3 \tan x - \tan^3 x}{1 - 3 \tan^2 x} \quad x \neq (2n+1)\frac{\pi}{6} \quad n \in \mathbb{Z} \quad (3)$$

(b) Hence solve, for  $0 < \theta < \frac{\pi}{2}$

$$\frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} = 2 \sec^2 3\theta - 8$$

giving your answers to 2 decimal places.

(5)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

## **Question 8 continued**



P 7 6 3 9 0 A 0 2 3 3 3 2

### **Question 8 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



## **Question 8 continued**

**(Total for Question 8 is 8 marks)**



9:


**Figure 2**

Figure 2 shows a sketch of the curve  $C$  with equation

$$x = \frac{2}{3} \sin\left(3y + \frac{\pi}{4}\right) \quad \frac{\pi}{12} < y < \frac{3\pi}{4}$$

The curve intersects the  $y$ -axis at the points  $A$  and  $B$  as shown.

(a) Find the exact value of the  $y$  coordinate of

- point  $A$
- point  $B$

(3)

(b) Show that

$$\left(\frac{dy}{dx}\right)^2 = \frac{1}{p - qx^2}$$

where  $p$  and  $q$  are integers to be found.

(4)

The **normal** to  $C$  at  $A$  and the **tangent** to  $C$  at  $B$  intersect at the point  $D$ .

Using

- the answer to part (b)
- the sketch of curve  $C$  in Figure 2

(c) find, in simplest form, the exact  $x$  coordinate of  $D$ .

(4)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

## **Question 9 continued**



P 7 6 3 9 0 A 0 2 7 3 2

### **Question 9 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



### **Question 9 continued**

(Total for Question 9 is 11 marks)




**10:****Figure 3**

Figure 3 shows a sketch of part of the graph with equation  $y = f(x)$ , where

$$f(x) = |kx - 10| + k \quad x \in \mathbb{R}$$

and  $k$  is a positive constant.

The graph

- cuts the  $y$ -axis at the point  $A$
- has a vertex at the point  $P$

(a) Find, in simplest form in terms of  $k$ ,

(i) the  $y$  coordinate of  $A$

(ii) the coordinates of  $P$

(3)

(b) Find, in terms of  $k$ , the range of values of  $x$  which satisfy

$$|kx - 10| + k \geq 2k$$

(3)

Given that the line with equation  $y = 3x + 1$  intersects the graph of  $y = f(x)$  at 2 distinct points,

(c) find the range of values of  $k$ .

(4)



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**Question 10 continued**



P 7 6 3 9 0 A 0 3 1 3 2

**Question 10 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**(Total for Question 10 is 10 marks)**

**TOTAL FOR PAPER IS 75 MARKS**

