

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

## Pearson Edexcel International Advanced Level

Monday 2 June 2025

Afternoon (Time: 1 hour 30 minutes)

Paper  
reference

**WME02/01**



### Mathematics

#### International Advanced Subsidiary/Advanced Level Mechanics M2

#### You must have:

Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

#### Instructions:

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need*.
- You should show sufficient working to make your methods clear.  
Answers without working may not gain full credit.
- Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ , and give your answer to either 2 significant figures or 3 significant figures.

#### Information:

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question*.

#### Advice:

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

*Turn over* ►

P76175A

©2025 Pearson Education Ltd.  
Y:1/1/1/1/



Pearson

1:

**In this question you must show all stages of your working.  
Solutions relying on calculator technology are not acceptable.**

A particle  $P$  moves along the  $x$ -axis.

At time  $t = 0$ ,  $P$  is at rest at the origin  $O$ .

At time  $t$  seconds,  $t \geq 0$ , the acceleration of  $P$ , in the positive  $x$  direction, is

$$k(3 - t^2) \text{ m s}^{-2}$$

where  $k$  is a positive constant.

Given that when  $t = 1.5$ , the speed of  $P$  is  $13.5 \text{ m s}^{-1}$

(a) show that  $k = 4$

(3)

The displacement of  $P$  from  $O$  is  $s$  metres in the positive  $x$  direction.

(b) Show that

$$s = \frac{1}{3} t^2 (18 - t^2)$$

(2)

(c) Find the total distance that  $P$  moves before it returns to  $O$ .

(4)



### Question 1 continued

**(Total for Question 1 is 9 marks)**



2:

**In this question you must show all stages of your working.  
Solutions relying on calculator technology are not acceptable.**

*[In this question,  $\mathbf{i}$  and  $\mathbf{j}$  are horizontal unit vectors.]*

A particle  $P$  of mass 3 kg is moving on a smooth horizontal plane.

At time  $t$  seconds, where  $1 \leq t \leq 4$ , the position vector,  $\mathbf{r}$  metres, of  $P$  relative to a fixed origin is given by  $\mathbf{r} = 4t^{\frac{1}{2}}\mathbf{i} - 3t\mathbf{j}$ .

At the instant when  $t = 4$ ,  $P$  receives an impulse  $(6\mathbf{i} - 3\mathbf{j})$  N s.

Find the speed of  $P$  immediately after the impulse is received.

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



## Question 2 continued

**(Total for Question 2 is 7 marks)**



A standard linear barcode is located at the top of the page, consisting of vertical black lines of varying widths on a white background. It is used for tracking and identification of the document.

3: [The distance of the centre of mass of a semicircular arc of radius  $r$  is  $\frac{2r}{\pi}$  from its centre.]




Figure 1

A uniform rigid wire **framework**, shown in Figure 1, consists of two wire semicircular arcs  $AB$  and  $CD$ , both with centre  $O$ , joined by two straight pieces of wire,  $AC$  and  $DB$ . Arc  $AB$  has radius  $2a$ , arc  $CD$  has radius  $a$ ,  $AC = DB = a$ , and  $ACDB$  is a straight line. The framework lies in a single plane.

(a) Show that the centre of mass of the framework is a distance  $\frac{6a}{(3\pi + 2)}$  from  $AB$ . (5)

The point  $E$  is on the arc  $AB$  and the angle  $AOE$  is a right angle.

The framework is freely suspended from  $A$ .

The framework hangs at rest in a vertical plane with  $AE$  making an angle  $\theta$  with the downward vertical.

(b) Find the exact value of  $\tan \theta$ , simplifying your answer. (5)



### Question 3 continued



### Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



### Question 3 continued

**(Total for Question 3 is 10 marks)**



A standard linear barcode is located at the top of the page, consisting of vertical black lines of varying widths on a white background.

4: A rough plane is inclined to the horizontal at an angle  $\alpha$ , where  $\sin \alpha = \frac{1}{7}$

A car of mass 750 kg moves up a line of greatest slope of the plane. The total resistance to the motion of the car from non-gravitational forces is 200 N. The engine of the car is working at a constant rate of 20 kW.

(a) Find the constant speed, in  $\text{ms}^{-1}$ , at which the car could move up the slope.

(4)

(b) Find the acceleration of the car at the instant when it is moving up the slope at  $10 \text{ m s}^{-1}$

(4)

At this instant, the car is at the point *A* and the engine of the car is switched off.

The total resistance to the motion of the car from non-gravitational forces remains at 200 N.

Using the work-energy principle,

(c) find how far up the slope from *A* the car would travel before coming to rest.

(4)



## Question 4 continued



## Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



## Question 4 continued

(Total for Question 4 is 12 marks)



5: A particle  $P$  of mass  $2m$  and a particle  $Q$  of mass  $m$  are at rest on a smooth horizontal plane. Particle  $P$  is projected with speed  $u$  along the plane towards  $Q$  and the particles collide. The coefficient of restitution between the particles is  $e$ , where  $e > 0$

(a) Show that the speed of  $Q$  after the collision is  $\frac{2(1+e)u}{3}$

(5)

The total loss of kinetic energy due to the collision is  $k(1 - e^2)mu^2$ , where  $k$  is a constant.

(b) Find the value of  $k$ .

(5)

The magnitude of the impulse of  $Q$  on  $P$  in the collision is  $\frac{8mu}{9}$

(c) Find the value of  $e$ .

(2)

After the collision,  $Q$  goes on to hit a vertical wall which is fixed at right angles to the direction of motion of  $Q$ . The coefficient of restitution between  $Q$  and the wall is  $f$ .

After  $Q$  has hit the wall,  $Q$  has the same speed as  $P$ .

(d) Find the value of  $f$ .

(3)



## Question 5 continued



## Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**DO NOT WRITE IN THIS AREA**



### Question 5 continued

**(Total for Question 5 is 15 marks)**



6:




Figure 2

A uniform rod  $AB$ , of mass  $m$  and length  $2a$ , has its end  $A$  on a rough horizontal floor.

A smooth horizontal rail is fixed above the floor.

The rod rests in equilibrium against the rail at the point  $C$ , where  $AC = 1.5a$ .

The vertical plane containing the rod is at right angles to the rail.

The rod is inclined to the floor at an angle  $\theta$ , as shown in Figure 2.

The magnitude of the normal reaction exerted on the rod at  $C$  is  $S$ .

(a) Show that  $S = \frac{2mg \cos \theta}{3}$  (3)

The magnitude of the vertical component of the force exerted on the rod at  $A$  is  $V$ .

(b) Show that  $V = \frac{mg}{3} (3 - 2 \cos^2 \theta)$  (3)

Given that the rod is in limiting equilibrium and the coefficient of friction between the rod and the floor is  $\frac{4}{7}$

(c) show that

$$p \tan^2 \theta + q \tan \theta + r = 0$$

where  $p$ ,  $q$  and  $r$  are integers to be found.

(6)



## Question 6 continued



## Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



## Question 6 continued

**(Total for Question 6 is 12 marks)**



7: A small stone is projected from a point  $O$  on horizontal ground.

The stone is projected with speed  $20\text{ m s}^{-1}$  at an angle  $\alpha$  to the ground.

When the stone is 11 m above the ground

- the vertical velocity component of the stone has magnitude  $V\text{ m s}^{-1}$
- the horizontal velocity component of the stone has magnitude  $2V\text{ m s}^{-1}$

In a model of the motion

- the stone is modelled as a particle moving freely under gravity
- the acceleration due to gravity is modelled as being  $10\text{ m s}^{-2}$**


Using the model and the principle of conservation of mechanical energy,

(a) show that  $V = 6$

(4)

Figure 3 shows a plane fixed to the ground at an angle of  $60^\circ$

The stone hits this plane at the point  $A$ .



**Figure 3**

The direction of the motion of the stone is perpendicular to the plane at the instant the stone hits  $A$ .

Using the model,

(b) find the height of  $A$  above the ground, giving your answer to 2 significant figures.

(6)



### Question 7 continued



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 7 is 10 marks)**

## **TOTAL FOR PAPER IS 75 MARKS**

