



Mark Scheme (Results)

Summer 2025

Pearson Edexcel International Advanced Level
In Pure Mathematics P2 (WMA12) Paper 01A

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2025

Question Paper Log Number P83172A

Publications Code WMA12_01A_2506_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Pearson Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for ‘knowing a method and attempting to apply it’, unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd or ft will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. All A marks are ‘correct answer only’ (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.
8. Marks for each question are scored by clicking in the marking grids that appear below each student response on ePEN. The maximum mark allocation for each question/part question(item) is set out in the marking grid and you should allocate a score of '0' or '1' for each mark, or "trait", as shown:

	0	1
aM		•
aA	•	
bM1		•
bA1	•	
bB	•	
bM2		•
bA2		•

9. Be careful when scoring a response that is either all correct or all incorrect. It is very easy to click down the '0' column when it was meant to be '1' and all correct.

General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use correct formula (with values for a , b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $(x \pm \frac{b}{2})^2 \pm q \pm c$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice

Question Number	Scheme					Marks
1.						B1 M1 A1 (3 marks)
	<i>a</i>	<i>b</i>	<i>c</i>	<i>abc</i>	One correct set <i>a, b & c</i>	
	2	15	3	90		
	3	9	8	216	Two correct rows	
	4	1	15	60		
	All values of <i>abc</i> are multiples of 6				Fully correct + statement	

Main method: Numerical answer finding all known solutions

B1: Any correct set of values for *a, b* and *c*. You can ignore any attempt at the product *abc*

M1: Any two fully correct rows including the correct calculations for the products.
Condone any extra or incorrect rows

A1: Fully correct. The solution must have

- the three correct rows with correct calculations for *abc*
- any extra (or incorrect) rows that may be present must be deleted or somehow excluded
- a minimal statement showing that they are aware that 90, 216 and 60 are multiples of 6.

E.g they are all divisible by 6, use of a tick ✓, proven, QED, calculations such as $\frac{90}{6} = 15$. They can be divided by 6 (on its own) is insufficient without further clarification such as "to give whole numbers"

- no incorrect calculations. E.g if their attempt to prove that 90 being a multiple of 6 includes $\frac{90}{6} = 10$ then it is A0

Algebraic solution or solutions that rely on explanations: These will be rare and unlikely to lead to completely correct solutions.

Example of an attempt that has some merit based upon odds and evens

If *a* is even

Then $a = 2n$, leading to $c = 4n^2 - 1 = (2n + 1)(2n - 1)$ and $b = 21 - 4n^2 - 2n$
giving $abc = 2n(21 - 4n^2 - 2n)(2n + 1)(2n - 1)$

If *a* is odd.

Then $a = 2n + 1$, leading to $c = 4n^2 + 4n = 4n(n + 1)$ and $b = 19 - 4n^2 - 6n$
giving $abc = (2n + 1)(19 - 4n^2 - 6n)4n(n + 1)$

B1: Attempts both odd and even and achieves correct algebraic expressions for *a, b* and *c* in both cases.

M1: Correct products with attempts when $n = 1, 2$ and achieves two fully correct rows

A1: Fully correct with no additional rows

Question Number	Scheme	Marks
2.(a)	E.g. $64 \times \left(-\frac{1}{2}\right)^3 = -8 \checkmark *$	M1, A1* (2)
(b)	Finds the value of a . E.g. $64 \div \left(-\frac{1}{2}\right)^2 = 256$ Uses $S_{\infty} = \frac{a}{1-r} = \frac{256}{1-\left(-\frac{1}{2}\right)} = \frac{512}{3}$	M1, A1 M1, A1 (4) (6 marks)

(a)

M1: Any acceptable attempt to prove the statement.

This usually involves getting to an equation in r ...

E.g. I $ar^2 = 64, ar^5 = -8 \Rightarrow 64r^3 = -8$ o.e. such as $ar^2 = 64, ar^5 = -8 \Rightarrow r^{-3} = -8$

E.g. II $ar^2 = 64, ar^5 = -8 \Rightarrow \frac{64}{r^2} = \frac{-8}{r^5}$

(the numbers used in the attempt, that is 64 and -8 must be correct)

but could involve using the given answer

E.g. III $64 \times \left(-\frac{1}{2}\right)^3, 64 \times -\frac{1}{2} \times -\frac{1}{2} \times -\frac{1}{2}$ or $-8 \div \left(-\frac{1}{2}\right)^3$

Condone the following types of slips/errors

E.g. I or II with incorrect indices on terms as long as they differ by 3;

$ar^3 = 64, ar^6 = -8 \Rightarrow r^3 = -\frac{8}{64}$

E.g. III with missing brackets $64 \times -\frac{1^3}{2}$

A1*: Fully correct proof. A minimal proof could be $r^3 = -\frac{8}{64} \Rightarrow r = -\frac{1}{2}$ o.e

As the result is given a direct proof must have a penultimate line equivalent of $r^3 = -\frac{8}{64}$ or $r^{-3} = -8$

If they state $64r^3 = -8$ followed by $r = -\frac{1}{2}$ without any intermediate work then it is M1 A0

If they assume the answer then expect to see equivalent to $64 \times \left(-\frac{1}{2}\right)^3 = -8$ which must be followed by a concluding statement (tick sufficient) is required.

Equivalent work would be $64 \xrightarrow{\times -\frac{1}{2}} -32 \xrightarrow{\times -\frac{1}{2}} 16 \xrightarrow{\times -\frac{1}{2}} -8 \checkmark$

This mark cannot be awarded from incorrect indices on r or missing brackets (that are necessary)

(b) **The marks for the following can only be awarded from work in part (b)**

M1: Uses a suitable method to find the value of a allowing $r = \pm \frac{1}{2}$

E.g. Attempts $a \times \left(\pm \frac{1}{2} \right)^2 = 64 \Rightarrow a = \dots$

Condone a missing bracket e.g. $a \times -\frac{1^2}{2} = 64 \Rightarrow a = \dots$

A1: States or implies that $a = 256$ via the correct $r = -\frac{1}{2}$.

Stating the value scores the two marks. Note that $a \times \frac{1}{4} = 64 \Rightarrow a = 256$ is M1, A1 (nothing incorrect)

This can be awarded following a missing bracket. (It isn't a proof)

M1: Uses $S_{\infty} = \frac{a}{1-r}$ with a value of $a \neq 64, -8$ and $r = \pm \frac{1}{2}$

So award $S_{\infty} = \frac{'a'}{1-\frac{1}{2}} = \dots$ or $S_{\infty} = \frac{'a'}{1-\left(-\frac{1}{2}\right)} = \dots$ but if an incorrect formula is stated score it M0

A1: $\frac{512}{3}$ o.e. ISW after a correct answer.

Condone $170.\dot{6}$ but do not allow awrt 170.7 unless the exact correct answer is seen first.

Question Number	Scheme	Marks
3 (a)	$f(x) = (3x^2 - 4x - 5)(x - k) - 5$ States -5	B1 (1)
(b)	Sets $f(-2) = 25 \rightarrow (3 \times 4 - 4 \times -2 - 5)(-2 - k) - 5 = 25$ $15(-2 - k) = 30 \Rightarrow -2 - k = 2 \Rightarrow k = -4$ *	M1 A1* (2)
(c)	$(3x^2 - 4x - 5)(x + 4) - 5 = 3x^3 + 8x^2 - 21x - 25$ Attempts $3x - 1 \overline{) 3x^3 + 8x^2 - 21x - 25}$ to achieve quotient of $\dots x^2 + \dots x + \dots$ and a remainder that is a constant Quotient = $x^2 + 3x - 6$ OR Remainder = -31 Quotient = $x^2 + 3x - 6$ AND Remainder = -31	B1 M1 A1ft A1 (4) (7 marks)

(a)

B1: States -5

You may see $f(k)$ or a division sum attempted, which is fine. You are just awarding the sight of -5

Don't be too concerned about what -5 is called. You may see $f(x)$, $f(k)$ or even x

(b)

M1: Sets $f(-2) = 25$ to form an equation in k Condone sign slips within the calculations but setting

$f(-2) = -25$ or $f(2) = 25$ is M0. Some may attempt to substitute -2 into the expanded form of

$f(x) = 3x^3 - (4 + 3k)x^2 + (4k - 5)x + 5k - 5$ which is fine. Condone slips but expect to see linear coefficients for k in each of x^2 , x and the constant terms.

Don't be concerned about poor attempts to solve the equation in k for this mark

A1*: Shows sufficient working with no errors to prove that $k = -4$.

The answer must follow a simplified and (easily) solvable equation.

(See scheme for possible route but allow simplified linear equations like $-30 - 15k = 30$)

.....
M1: Alternatively substitutes $k = -4$ into $f(x)$ to get $(3x^2 - 4x - 5)(x + 4) - 5$ and attempt the value of $f(-2)$

A1: Shows that $f(-2) = (12 + 8 - 5)(-2 + 4) - 5 = 15 \times 2 - 5 = 25$ and makes a minimal conclusion e.g. ✓

.....

(c)

B1: Multiplies out $(3x^2 - 4x - 5)(x + 4) - 5$ to achieve a simplified $3x^3 + 8x^2 - 21x - 25$

M1: Scored for a full method of finding **both** the quotient and the remainder.

The two main ways of scoring this mark are;

Either attempting to divide their expanded $f(x)$ by $(3x - 1)$ to form a quotient of the form

$Ax^2 + B.x + C$ and a remainder that is a constant.

Alternatively, setting up an identity of the form $(Ax^2 + B.x + C)(3x - 1) + R$ followed by an attempt to find the values of A, B, C and R .

Watch for candidates who find $f\left(\frac{1}{3}\right) = -31$ and then adapt $f(x)$ to $3x^3 + 8x^2 - 21x + 6$. The M mark

would be scored for dividing their $3x^3 + 8x^2 - 21x + 6$ by $(3x - 1)$ to form a quotient of the form

$Ax^2 + B.x + C$ and a remainder that is a constant (which should be 0 in a correct division).

A1ft: For either the quotient = $x^2 + 3x - 6$. Allow this to be scored within the division sum

Or the remainder = -31 (may be scored within the division sum)

It must follow M1 so attempting $f\left(\frac{1}{3}\right)$ and reaching -31 will not score this mark without further

work

Follow through on finding a correct quotient or remainder for their $3x^3 + 8x^2 - 21x - 25$ (4 term cubic expression). Note that any $3x^3 + 8x^2 - 21x + k$ will have a quotient of $x^2 + 3x - 6$

A1: States that the Quotient = $x^2 + 3x - 6$ AND the Remainder = -31 . These cannot be scored within a division sum and must be stated separately to this. All previous marks must have been scored in (c) so cannot be awarded following an incorrect $f(x)$

Question Number	Scheme	Marks
4. (a)	$\log_3(a+1) - \log_3 a = 4 \Rightarrow \log_3\left(\frac{a+1}{a}\right) = 4$ $\Rightarrow \left(\frac{a+1}{a}\right) = 3^4$ $\Rightarrow a+1 = 81a \Rightarrow a = \frac{1}{80}$	M1 A1 dM1, A1 (4)
(b)	$\text{Area} \approx \frac{1}{2} \left\{ \log_3\left(\frac{2}{1}\right) + \log_3\left(\frac{6}{5}\right) + 2 \left(\log_3\left(\frac{3}{2}\right) + \log_3\left(\frac{4}{3}\right) + \log_3\left(\frac{5}{4}\right) \right) \right\}$ $= \frac{1}{2} \log_3 \left(\frac{2}{1} \times \frac{6}{5} \times \frac{3^2}{2^2} \times \frac{4^2}{3^2} \times \frac{5^2}{4^2} \right) = \frac{1}{2} \log_3 15 = \log_3 \sqrt{15}$	M1, A1 dM1, A1 (4)
(c)	States 'increase the number of strips'	B1 (1) (9 marks)

(a) Allow with $a \leftrightarrow x$ as seen in the notes

M1: Sets $\log_3(x+1) - \log_3 x = 4$ and then applies the subtraction law. Condone the omission of the 3 in \log_3

Alternatively writes the equation $\log_3(x+1) = 4 + \log_3 x$ as $\log_3(x+1) = \log_3 k + \log_3 x$ ($k \neq 4$)

and then uses the addition law $\log_3(x+1) = \log_3(kx)$

A1: Correct equation not involving logs. For example, $\left(\frac{x+1}{x}\right) = 3^4$

dM1: Correct attempt to solve an equation of the form $\frac{x+1}{x} = k$ o.e. where k is an integer $k \neq 3$ or 4.

Look for $\frac{x+1}{x} = k \Rightarrow x = \frac{1}{k-1}$ where $k \neq 3$.

A1: a or $x = \frac{1}{80}$ or 0.0125 following the award of M1, A1, dM1.

Working must be seen as required by the question.

Special Case:

You may see attempts that proceed towards the correct answer using incorrect laws.

E.g. $\log_3(x+1) - \log_3 x = 4 \Rightarrow \frac{\log_3(x+1)}{\log_3 x} = 4 \Rightarrow \left(\frac{x+1}{x}\right) = 3^4 \Rightarrow x = \frac{1}{80}$

Candidates who achieve the correct answer using the above incorrect line can score SC 0, 0, 1, 1

(b)

Condone the omission of the 3 in \log_3 except for the final A mark where they must arrive at $\log_3 \sqrt{15}$.

Any use of calculators to produce decimal equivalents to these values will result in a maximum score of 1,0,0,0 marks as an exact value could never be achieved.

M1: Score for a reasonable attempt at applying the trapezium rule.

Look for **an attempt** at the correct form $\frac{h}{2} \{ \text{first term} + \text{last term} + 2(\text{sum of the other terms}) \}$ with at least the following

- (1) Correct first **or** last term
- (2) At least two correct 'other terms'
- (3) $h = 1$

The bracketing must be correct but condone a missing trailing bracket.

So, look for a minimum of $\frac{1}{2} \{ a + e + 2 \times (b + c + d) \}$ with a or e correct, two out of three of b , c and d correct with the missing trailing bracket

FYI the decimal solution is $\frac{1}{2} \{ 0.631 + 0.166 + 2(0.369 + 0.262 + 0.203) \}$ allow awrt 2 dp

A1: A fully correct application of the trapezium rule in un-simplified form.

Amongst various correct alternatives are

$$\text{Area} \approx \frac{1}{2} \left\{ \log_3 \left(\frac{2}{1} \right) + \log_3 \left(\frac{6}{5} \right) + 2 \left(\log_3 \left(\frac{3}{2} \right) + \log_3 \left(\frac{4}{3} \right) + \log_3 \left(\frac{5}{4} \right) \right) \right\}$$

$$\text{Area} \approx \frac{1}{2} \{ \log_3 2 + \log_3 6 - \log_3 5 + 2(\log_3 3 - \log_3 2 + \log_3 4 - \log_3 3 + \log_3 5 - \log_3 4) \}$$

$$\text{Area} \approx \frac{1}{2} \{ \log_3 2 + \log_3 6 - \log_3 5 + 2(-\log_3 2 + \log_3 5) \}$$

$$\text{Area} \approx \frac{1}{2} \left\{ \log_3 2 + \log_3 \frac{3}{2} \right\} + \frac{1}{2} \left\{ \log_3 \frac{3}{2} + \log_3 \frac{4}{3} \right\} + \frac{1}{2} \left\{ \log_3 \frac{4}{3} + \log_3 \frac{5}{4} \right\} + \frac{1}{2} \left\{ \log_3 \frac{5}{4} + \log_3 \frac{6}{5} \right\}$$

FYI: A fully simplified expression for the approximate area is $\frac{1}{2} \{ \log_3 6 + \log_3 5 - \log_3 2 \}$

Condone the omission of the trailing bracket as long as subsequent work implies its presence.

dM1: Correct use of logs in an attempt to fully combine their terms following the award of the previous M.

This must include sight of at least one correct use of two of the three log laws.

So, use of just addition and subtraction laws is fine as would addition and power laws

This mark can be implied if they go directly from a correct exact expression to the correct answer.

Candidates must proceed as far as $\frac{1}{2} \log p$ or $\log q$

A1: $\log_3 \sqrt{15}$ following M1, A1 dM1

(c)

B1: States

- 'increase the number of strips'
- 'decrease the width of the strips'
- 'make more strips'

without any incorrect statements such as increase the strips, so try $\int_1^{20} \log_3 (x+1) - \log_3 x \, dx$

Do not allow vague statements such as 'add more numbers of x '

Question Number	Scheme	Marks
5. (a)	$u_2 = 1 - \frac{1}{4} = \frac{3}{4}, \quad u_3 = 1 - \frac{4}{3} = -\frac{1}{3},$ $u_4 = 1 - \left(-\frac{3}{1}\right) = 4 \quad \text{with statement saying the sequence repeats every 3 terms o.e}$	M1 A1 A1* (3)
(b)	$\sum_{n=1}^{180} (5n+3) = \frac{180}{2} (8+903) = (81990)$ $\sum_{n=1}^{180} (u_n) = 60 \times \left(4 + \frac{3}{4} + -\frac{1}{3}\right) = (265)$ $\sum_{n=1}^{180} (5n+3+u_n) = 81990 + 265 = 82255$	M1 M1 A1, A1 (4) (7 marks)

(a)

M1: Attempts to use the iterative sequence correctly at least once.

Most often this is for $u_2 = 1 - \frac{1}{4}$ or $u_2 = \frac{3}{4}$ o.e.

Condone omission of the lhs as long as it is obvious. E.g. 4, 0.75

A1: Achieves a correct value for both u_2 and u_3 . Condone $u_3 = -0.\dot{3}$

Do not allow $u_3 = -0.33$ unless you see the exact value beforehand.

Allow $(4), \frac{3}{4}, -\frac{1}{3}$ if it is obviously values for both u_2 and u_3 .

A1*: This is a show that question and requires

- correct values for u_2 and u_3
- a correct calculation and value seen for u_4 E.g. $u_4 = 1 - \left(-\frac{3}{1}\right) = 4$
- a minimal reason/statement 'the sequence repeats every 3 terms' o.e such as 'as $u_4 = u_1$,
- a minimal conclusion such as sequence is periodic of order 3' or 'hence true'

(b) **All stages of working must be shown so don't accept just answers as evidence of method**

M1: Establishes a viable method to find $\sum_{n=1}^{180} (5n+3)$ using the fact that it is an AP.

Look for use of $\frac{n}{2}\{a+l\}$ or $\frac{n}{2}\{2a+(n-1)d\}$ o.e. with $n=180, a=3, 5 \text{ or } 8$ and $d=5$

Alternatively attempts $\sum_{n=1}^{180} (5n+3) = \sum_{n=1}^{180} 5n + \sum_{n=1}^{180} 3 = 5 \times \frac{180}{2} \{180+1\} + 180 \times 3$ o.e.

Do not just accept $\sum_{n=1}^{180} (5n+3) = 81990$ without sight of an intermediate calculation

This may be awarded from terms within a more complicated expression to complete $\sum_{n=1}^{180} (5n+3+u_n)$

M1: Establishes a viable method to find $\sum_{n=1}^{180} (u_n)$.

For example, as it repeats every 3 terms look for $60 \times (4+u_2+u_3)$ o.e such as

$$60 \times 4 + 60 \times u_2 + 60 \times u_3$$

Do not accept $\sum_{n=1}^{180} (u_n) = 265$ without sight of an intermediate calculation

This may be awarded from terms within a more complicated expression to complete $\sum_{n=1}^{180} (5n+3+u_n)$

A1: Scored for either $\sum_{n=1}^{180} (5n+3) = 81990$ or $\sum_{n=1}^{180} (u_n) = 265$ following the award of the equivalent M mark

It can be implied, so for instance 82255 following the award of both M's but it is the value, not the calculation that must be seen or implied.

A less obvious award would be for $\sum_{n=1}^{180} (u_n+3) = 60 \times \left(4 + \frac{3}{4} + -\frac{1}{3} + 9\right) = 805$ which implies the 265

A1: 82255 which must follow the award of M1, M1 and A1

Question Number	Scheme	Marks
6 (i) (a)	Uses $\sin^2 \theta + \cos^2 \theta = 1$ with $\cos \theta = \frac{1}{\sqrt{5}} \Rightarrow \sin^2 \theta = \frac{4}{5}$	M1
	$\Rightarrow \sin \theta = -\frac{2}{\sqrt{5}}$	A1
	(b) Uses $\tan \theta = \frac{\sin \theta}{\cos \theta}$ with $\cos \theta = \frac{1}{\sqrt{5}}$ and their $\sin \theta = -\frac{2}{\sqrt{5}}$	M1
	$\Rightarrow \tan \theta = -2$	A1
		(4)
(ii) (a)	1 (m)	B1
		(1)
(b)	$30t - 40 = 180 \Rightarrow t = 7.33$ Hence 7:20 am	M1, A1, A1
		(3)
(c)	$4 + 3 \cos(30T - 40)^\circ = 3.5 \Rightarrow \cos(30T - 40)^\circ = -\frac{1}{6}$	M1
	$\Rightarrow 30T - 40 = 99.6, 260.4, \underline{459.6}$	A1
	$\Rightarrow T = \frac{"459.6" + 40}{30} = 16.65$	dM1, A1
		(4)
		(12 marks)

(i)

Clear use of trigonometric identities must be shown. Answers without method don't score marks



Look for work within the body of the script which may be common

(a)

M1: Uses $\sin^2 \theta + \cos^2 \theta = 1$ with $\cos \theta = \frac{1}{\sqrt{5}} \Rightarrow \sin^2 \theta + \frac{1}{5} = 1$ o.e.

Alternatively shows a 'triangle' method and uses Pythagoras' theorem to reach $\sin \theta = \pm \frac{2}{\sqrt{5}}$

E.g.



A1: $\sin \theta = -\frac{2}{\sqrt{5}}$ or simplified equivalent such as $-\frac{2\sqrt{5}}{5}$ following the award of the M1. ISW after a

correct answer. Note that $\sin \theta = -\sqrt{\frac{4}{5}}$ is A0 as it is not simplified

(b)

M1: Uses $\tan \theta = \frac{\sin \theta}{\cos \theta}$ with $\cos \theta = \frac{1}{\sqrt{5}}$ and either $\sin \theta = \frac{2}{\sqrt{5}}$ or $\sin \theta = -\frac{2}{\sqrt{5}}$ from (a) which must have been awarded from a correct method.

Alternatively, from a triangle method, shows the opposite side = 2 in a right-angled triangle and takes the correct ratio

A1: $\tan \theta = -2$ following the award of the M1 in both (i) and (ii)

(ii)(a)

B1: 1 (m) Units are not necessary here but incorrect units e.g. 1 cm would be B0

(ii)(b)

M1: Sets $30t - 40$ equal to any value in degrees where the cosine is -1

So $30t - 40 = 180(2n+1)$, for any $n \in \mathbb{Z}$ is acceptable. Condone $30t - 40 = (2n+1)\pi$,

Condone the result obtained from differentiation with $\sin(30t - 40) = 0 \Rightarrow 30t - 40 = 180(2n+1)$

A1: $30t - 40 = 180 \Rightarrow t = 7\frac{1}{3}$ or 7.33 but condone 7.3. It must follow the award of the M mark

A1: Achieves 7:20 am or 07:20 or any other acceptable way of establishing the exact **time of day** time in hours and minutes. So 7 hours 20 minutes after midnight is fine. **Condone 7:20 but 7:20pm is A0 and 7 hours 20 minutes without reference to after midnight is A0. It must follow correct working and cannot be rounded from a different time value such as 7:19.5**

(ii)(c) Radian attempts can only usually score the first M1.

M1: Sets $4 + 3\cos(30T - 40)^\circ$ equal to 3.5 and proceeds to $\cos(30T - 40)^\circ = k$, $|k| < 1$

A1: Achieves $30T - 40$ equal to any of 99.6, 260.4, 459.6

dM1: Proceeds from $30T - 40$ equal to any of awrt 100, 260, 460 AND uses correct order of operations and working leading to a value for T .

For example, allow for $30T - 40 = \text{awrt } 100 \Rightarrow T = \frac{140}{30}$ or awrt 4.7

$$30T - 40 = \text{awrt } 260 \Rightarrow T = \text{awrt } 10$$

$$30T - 40 = \text{awrt } 460 \Rightarrow T = \frac{500}{30} \text{ or awrt } 16.7$$

A1: awrt 16.65 following the award of M1, A1 dM1

ISW after a correct answer for T . Ignore any attempts to convert the T value to a time.

A clear method must be shown here, so solutions produced entirely from a calculator will not score.

E.g. $4 + 3\cos(30T - 40)^\circ = 3.5 \Rightarrow T = 16.65$ is 0 marks

A minimal acceptable response for 4 marks would be

$$4 + 3\cos(30T - 40)^\circ = 3.5 \Rightarrow \cos(30T - 40)^\circ = -\frac{1}{6}$$

$$\Rightarrow 30T - 40 = 459.6$$

$$\Rightarrow T = 16.65$$

Question Number	Scheme	Marks
7 (a)	States either $kn = -24$ (1) or $\frac{n(n-1)}{2}k^2 = 270$ (2)	M1
	States both $kn = -24$ (1) and $\frac{n(n-1)}{2}k^2 = 270$ (2)	A1
	Substitutes $k = -\frac{24}{n}$ in equation (2) $\Rightarrow \frac{n(n-1)}{2} \left(-\frac{24}{n}\right)^2 = 270 \Rightarrow n = \dots$	M1
	$n = 16$	A1
	Uses their $n = 16$ in $kn = -24 \Rightarrow k = -\frac{24}{16} = -\frac{3}{2}$	dM1, A1
		(6)
(b)	$p = \frac{n(n-1)(n-2)}{3!}k^3 = \frac{16 \times 15 \times 14}{6} \times \left(-\frac{3}{2}\right)^3 = \dots$	M1
	$= -1890$	A1
		(2)
		(8 marks)

(a)

M1: States either $kn = -24$ (1) or $\frac{n(n-1)}{2}k^2 = 270$ (2) accepting equivalents but see note below

This mark cannot be awarded from $\binom{n}{1}k = -24$ or ${}^nC_2(kx)^2 = 270x^2$, with factorials

$$\frac{n!}{(n-2)!}k^2 = 540$$

or indeed versions including x terms such as $kx = -24x$ unless followed by a correct $kn = -24$

A1: States both $kn = -24$ (1) and $\frac{n(n-1)}{2}k^2 = 270$ (2) accepting equivalents

M1: Forms an equation in one variable leading to a value for k or n

It is dependent upon having achieved $kn = \pm 24$ and $\frac{n(n-1)}{2}k^2 = 270$

The process of forming the equation must be credible so look for $k = \frac{\pm 24}{n}$ being substituted and

$$\text{NOT } k = \pm 24 - n \quad \text{FYI the equation in } k \text{ is } \Rightarrow -\frac{24}{k} \left(-\frac{24}{k} - 1\right) \times \frac{k^2}{2} = 270 \Rightarrow k = \dots$$

A1: For either $n = 16$ following a correctly solved equation in n .

or $k = -\frac{3}{2}$ if the equation in k was formed.

Look for correct working leading to the value for n . For example, forming a linear equation is

$$\text{sufficient. } \frac{n(n-1)}{2} \left(-\frac{24}{n}\right)^2 = 270 \Rightarrow \frac{288(n-1)}{n} = 270 \Rightarrow 288n - 288 = 270n \Rightarrow n = 16$$

dM1: Uses $k = \pm \frac{24}{n}$ o.e. with their $n = '16'$ to find k or their $k = '-\frac{3}{2}'$ to find n

This mark may be implied. It is dependent upon the **previous M**

So, if they have scored the previous M, achieved $n = 8$ and $k = \frac{24}{n}$ award for $k = 3$

A1: $n = 16$ and $k = -\frac{3}{2}$ following the award of all previous marks

.....
The demand of the question is that they show all stages of their working.

Example I

If a candidate states both equations $kn = -24$, $\frac{n(n-1)}{2}k^2 = 270$, shows **a correct intermediate line in a single variable** then follows this with correct $n = 16$ and $k = -\frac{3}{2}$ **without the necessary working** they can be awarded SC: 1, 1, 1, 0, 1, 0

Example II

If a candidate states both equations $kn = -24$, $\frac{n(n-1)}{2}k^2 = 270$ and follows this with correct $n = 16$ and $k = -\frac{3}{2}$ without any necessary intermediate equations and working they can be awarded SC: 1, 1, 0, 1, 0, 0

If the SC is seen in part (a), they can go on to score both marks in (b) for correct answers

.....
(b)

M1: Uses $p = \frac{n(n-1)(n-2)}{3!}k^3$ with their values for n and k found from part (a) to find a value for p .

This can be implied by a 'correct' answer for their values of n and k .

Allow fractional values for both n and k

Allow this to be scored by candidates who perhaps have only one equation, usually $kn = -24$ and go on to state for example that $n = 4$ and $k = -6$ and then find $p = -864$

Values cannot be made up without an attempt in part (a) scoring at least one mark

A1: -1890 following correct values for n and k .

It must follow correct equations in part (a) but allow the values to be found from equations containing terms including factorial and/or combination notation.

Question Number	Scheme	Marks
8. (a)	$-5 < x < \frac{2}{3}$	M1, A1 (2)
(b)	$(2x-7)$	B1 (1)
(c)	$f'(x) = 2(3x-2)(x+5) = 6x^2 + 26x - 20$ $f(x) = 2x^3 + 13x^2 - 20x + c$ $x = \frac{7}{2}, \quad y/f(x) = 0 \Rightarrow c = (-175)$ $f(x) = 2x^3 + 13x^2 - 20x - 175 = (2x-7)(x^2 + 10x + 25)$ $= (2x-7)(x+5)^2$	M1, A1 dM1, A1 ddM1 A1 (6) (9 marks)

(a)

M1: Achieves the correct critical values. Condone for this mark -5 and awrt 0.67

They may be seen as part of an incorrect inequality

A1: $-5 < x < \frac{2}{3}$ or $-5, x, \frac{2}{3}$. Either of these on their own (with consistent inequality use) scores both marks

Accept other valid alternatives such as $x \in \left(-5, \frac{2}{3}\right)$ and $\{x : x > -5\} \cap \left\{x : x < \frac{2}{3}\right\}$ but not $x < \frac{2}{3}$ or $x > -5$

(b)

B1: States $(2x-7)$ or $A = 2, B = -7$ which must be seen in (b).

Condone candidates who write $\left(x - \frac{7}{2}\right) = (2x-7)$ Note that stating $\left(x - \frac{7}{2}\right)$ on its own is B0

(c) Some candidates integrate in part (b). Allow these candidates to score the marks in (c) as long as they re-write $f(x)$ in (c) or do some extra work involving $f(x)$ in (c)

M1: Multiplies out $f'(x)$ to form a quadratic expression and attempts to integrate.

Look for $f'(x) = 2(3x-2)(x+5) = ax^2 + bx + c \rightarrow f(x) = px^3 + qx^2 + rx \dots (+c)$ with or without the +
c

A1: $f(x) = 2x^3 + 13x^2 - 20x + c$ with or without the + c

dM1: Substitutes $x = \frac{7}{2}, \quad y/f(x) = 0 \Rightarrow c = \dots$

Alternatives exist via division using the fact that $(2x-7)$ is a factor of $2x^3 + 13x^2 - 20x + c$

A1: $f(x) = 2x^3 + 13x^2 - 20x - 175$ but allow $f(x) = 2x^3 + 13x^2 - 20x + c$ followed by $c = -175$

ddM1: Attempts to divide or factorise out a factor of $(2x-7)$ or $\left(x-\frac{7}{2}\right)$ from their

$$f(x) = 2x^3 + 13x^2 - 20x - 175$$

Awarded for 'correct' first and last terms or correct first two terms for their $f(x)$ condoning sign slips

$$\text{E.g. } 2x^3 + 13x^2 - 20x - 175 = (2x-7)(x^2 + \dots x \pm 25) \text{ or}$$

$$2x^3 + 13x^2 - 20x - 175 = \left(x - \frac{7}{2}\right)(2x^2 + \dots x \pm 50)$$

It is dependent upon both previous M's so cannot be awarded if they don't have a value for c
Mark their final response here as there may be working before the final answer

A1: CSO Requires working to be seen (see bold sentences in question) so look for

- $f(x) = (2x-7)(x+5)^2$ or $f(x) = (2x-7)(x+5)(x+5)$ following correct $f(x)$
- AND correct intermediate factorisation $(2x-7)(x^2 + 10x + 25)$

Going directly from $2x^3 + 13x^2 - 20x - 175$ to $(2x-7)(x+5)^2$ without sight of working would result in the loss of the last two marks

Note that the last 4 marks can be scored without finding the value of c BUT the constant c must be present.

We know that $f(x) = 2x^3 + 13x^2 - 20x + c$ and that $(2x-7)$ is a factor

So $2x^3 + 13x^2 - 20x + c = (2x-7)(Ax^2 + Bx + C)$ which may be attempted via division

$$2x^3 + 13x^2 - 20x + c = (2x-7)(1x^2 + 10x + 25) = (2x-7)(x+5)^2 \text{ o.e.}$$

The 6 marks in part (c) are awarded as follows

M1, A1: For $f(x) = 2x^3 + 13x^2 - 20x + c$ with or without the $+c$


dM1: For setting their $f(x) = '2'x^3 + '13'x^2 - '20'x + c = (2x-7)(Ax^2 + Bx + C)$ and correctly deduces the value for A for their $f(x)$. You may see a division attempted here which is equivalent work

A1: Correctly deduces $2x^3 + 13x^2 - 20x + c = (2x-7)(x^2 + 10x + C)$

Via division, the $(x^2 + 10x + \dots)$ can be awarded implied from within the calculation

ddM1: Correctly deduces $2x^3 + 13x^2 - 20x + c = (2x-7)(x^2 + 10x + 25)$

A1: $f(x) = (2x-7)(x+5)^2$ o.e.

Question Number	Scheme	Marks
9(a)		B1
(b)	$(x \pm a)^2 + y^2 = \dots$ $(x \pm a)^2 + y^2 = a^2$ <p>Uses (5, 6) in $(x \pm a)^2 + y^2 = a^2$ to form and solve an equation in a</p> <p>E.g. $(5 - a)^2 + 36 = a^2 \Rightarrow 10a = 61 \Rightarrow a = 6.1$</p> $(x - 6.1)^2 + y^2 = 6.1^2$	M1 A1 dM1 A1 (4) (5 marks)

(a)

B1: A circle lying in quadrants 1 and 4 only which

- passes through (0, 0)
- is symmetrical about the x -axis

Condone slips of the pen. Mark this positively.

Ignore any reference to the point (5, 6)

Ignore any reference to the coordinates of the centre

(b) Way One: Candidates start from an appropriate equation involving x^2 and y^2 then use (5, 6)

The following equations are mostly written with a constant a . Any suitable constant may be used.

The sign of the constant is also not important so accept $(x + a)^2 + y^2 = \dots$ or $(x - a)^2 + y^2 = \dots$

as well as $(a - x)^2 + y^2 = \dots$

M1: Allowable left-hand side for the equation of C For example $(x \pm a)^2 + y^2 = \dots$ or $(a - x)^2 + y^2 = \dots$

This must involve a constant and not be $(x - 5)^2 + y^2 = \dots$ for example

A1: Correct form for the equation involving a **single** constant.

Examples of correct equations are $(x - a)^2 + y^2 = a^2$, $(x + k)^2 + y^2 = k^2$ and $x^2 + y^2 = \pm 2px$

If they write $(x - a)^2 + y^2 = r^2$ and $r^2 = a^2$ this can be implied

dM1: Uses (5, 6) in $(x \pm a)^2 + y^2 = a^2$ o.e. to form and solve an equation in a .


E.g. $(5 - a)^2 + 36 = a^2 \Rightarrow \pm 10a = 61 \Rightarrow a = \dots$

$5^2 + 6^2 = \pm 10p \Rightarrow p = \dots$

A1: $(x - 6.1)^2 + y^2 = 6.1^2$ o.e. such as $x^2 + y^2 = 12.2x$

.....

Way Two: Candidates use Pythagoras' theorem to set up an equation in single variable



M1: Applies Pythagoras's theorem leading to the equation $(r-5)^2 + 36 = r^2$ or $(5-r)^2 + 36 = r^2$

This may be awarded for any variable

A1: Solves the above equation to find correct value $r = 6.1$

dM1: Uses a correct equation of circle $(x - '6.1')^2 + y^2 = '6.1'^2$ following through on their 6.1.

This is dependent on the 6.1 being a solution of a correct equation (previous M1)

A1: $(x - 6.1)^2 + y^2 = 6.1^2$ o.e.

Way Three: Bisector of chord method


M1: Attempts to find where the perpendicular bisector cuts the x -axis
Look for a good attempt using mid points and perpendicular gradient work.

A1: Achieves equation of perpendicular bisector as $y = -\frac{5}{6}x + \frac{61}{12}$

and point of intersection with the x -axis to be $\frac{61}{10}$ o.e

dM1: Correct form of equation with $(x - '6.1')^2 + y^2 = '6.1'^2$

A1: $(x - 6.1)^2 + y^2 = 6.1^2$ o.e.



Question Number	Scheme	Marks
10 (a)	$y = 2x + \frac{64}{x^2} - 3$ $\frac{dy}{dx} = 2 - \frac{128}{x^3}$ <p>Attempts to solve $\frac{dy}{dx} \Rightarrow x^3 = 64 \Rightarrow x = 4$ *</p>	<p>M1, A1</p> <p>dM1, A1*</p> <p>(4)</p>
(b)	$\int 2x + \frac{64}{x^2} - 3 \, dx = x^2 - \frac{64}{x} - 3x \quad (+c)$ <p>Finds the y values at both $x = 2$ and $x = 4$. $M = (0, 17)$ and $N = (0, 9)$</p> <p>Full attempt at area =</p> $\left[x^2 - \frac{64}{x} - 3x \right]_2^4 - 9 \times 2 + 2 \times (17 - 9) = 22 - 18 + 16 = 20$	<p>M1, A1</p> <p>B1</p> <p>dM1, A1</p> <p>(5)</p> <p>(9 marks)</p>

(a)

M1: Attempts $\frac{dy}{dx}$ with one correct aspect. Allow for $2x \rightarrow 2$ or $x^{-2} \rightarrow x^{-3}$ but not $3 \rightarrow 0$

A1: $\frac{dy}{dx} = 2 - \frac{128}{x^3}$ o.e which may be left un simplified. The lhs is not required

dM1: Sets their $\frac{dy}{dx} = p \pm qx^{-3} = 0$ and proceeds to $\alpha x^{\pm 3} = \beta$

Also allow candidates to substitute 4 into their $\frac{dy}{dx}$ which must be of the form $\frac{dy}{dx} = p \pm qx^{-3}$ and find its value.

A1*: Shows that the x coordinate of P is 4. This is a given answer so there must be sufficient evidence

For a direct proof you should see all of $2 - \frac{128}{x^3} = 0$, $x^3 = 64$ (condone $2x^3 = 128$) and $x = 4$

or $2 - \frac{128}{x^3} = 0$, $x^{-3} = \frac{1}{64}$ (condone $x^{-3} = \frac{2}{128}$) and $x = 4$

For a substitution method look for $\frac{dy}{dx} = 2 - \frac{128}{x^3}$ and at P, $\frac{dy}{dx} = 2 - \frac{128}{4^3} = 0$, hence proven

(o.e.)

(b)

M1: Attempts to integrate $2x + \frac{64}{x^2} - 3$ scored for increasing a correct index by 1 including $3 \rightarrow 3x$

A1: $\int 2x + \frac{64}{x^2} - 3 \, dx = x^2 - \frac{64}{x} - 3x \quad (+c)$ which may be left un simplified

B1: Finds the y values at both $x = 2$ and $x = 4$. $M = (0, 17)$ and $N = (0, 9)$

This mark may be implied by an area calculation $\left[x^2 - \frac{64}{x} - 3x \right]_2^4 - \underline{9} \times 2 + 2 \times \underline{8}$

Evidence may be awarded from the diagram or from coordinates for P and Q

dM1: Full attempt at area. It dependent upon having done some correct integration (M1 scored).

Look for the 'correct' areas being combined in the right way, condoning slips in integration and arithmetical slips

Total area = '22' - '18' + '16' = ...

A1: 20

Alt method to b using (curve – line) can be marked in a similar way

M1: Attempts to integrate $\left(2x + \frac{64}{x^2} - 3 \right) - (k)$ scored for increasing a correct index by 1

A1: $\int 2x + \frac{64}{x^2} - 12 \, dx = x^2 - \frac{64}{x} - 12x \quad (+c)$ which may be left un simplified

B1: Finds the y values at both $x = 2$ and $x = 4$. $M = (0, 17)$ and $N = (0, 9)$

This mark may be implied by an area calculation $\left[x^2 - \frac{64}{x} - 12x \right]_2^4 + (\underline{17} - \underline{9}) \times 2$

dM1: Full attempt at area. It dependent upon having done some correct integration (M1 scored).

Shaded area under curve $R_1 = \int_2^4 \left(2x + \frac{64}{x^2} - 3 \right) - (9) \, dx = \left[x^2 - \frac{64}{x} - 12x \right]_2^4 = 4$

Total area = $4 + 16 = 20$

Look for the 'correct' areas being combined in the right way, condoning slips in integration and arithmetical slips (such as $4 + 16 = 22$)

A1: 20

