



Mark Scheme (Results)

October 2025

International Advanced Level in Statistics S1

WST01/01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2025

Question Paper Log Number P84842A

Publication Code WST01_01_2510_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for ‘knowing a method and attempting to apply it’, unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes.

 - bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are ‘correct answer only’ (cao.), unless shown, for example, as A1ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer

Question Number	Scheme		Marks
1(a)	$[S_{hw}] = 604135 - \frac{1160 \times 4160}{8}$ or $[S_{ww}] = 2166600 - \frac{4160^2}{8}$ $[S_{hw}] = 935$ $[S_{ww}] = 3400$		M1 A1 A1 (3)
(b)	$r = \frac{"935"}{\sqrt{286 \times "3400"}}$ = 0.9481... awrt 0.948		M1A1 (2)
(c)	$b = \frac{"935"}{286} [= 3.269...]$ $a = \frac{4160}{8} - "3.269..." \times \frac{1160}{8} [= 45.96...]$ $w = 46.0 + 3.27h^*$		M1 M1 A1* (3)
(d)	[On average] for each/1 [cm] increase in height/h the weight/w increases by “3.27” [kg]		B1 (1)
(e)(i)	546.31 awrt 546		B1
(ii)	601.9 awrt 602		B1 (2)
(f)	The 1 st estimate is more reliable estimate since... • The 1 st estimate is an interpolation /The 2 nd is an extrapolation • 153(height) is within data range/170 is not within data range • 153 is closer to the mean(height)/170 is further from the mean		B1 (1)
	Notes		Total 12
(a)	M1	for a correct expression for S_{hw} or S_{ww} implied by either correct answer	
	A1	935 cao	
	A1	3400 cao	
(b)	M1	for a valid attempt at r allow ft their values from (a)	
	A1	awrt 0.948 correct answer scores 2 out of 2	
(c)	M1	for a correct numerical expression to find the value of b	
	M1	for a correct method to find the value of a ft their b Allow $w - 520 = "3.269..."(h - 145)$ oe for this mark	
	A1*	correct solution with both method marks and correct given equation stated. do not award this mark if there is incorrect working seen e.g. $a =$	

		45.85 rounding to $a = 46.0$ do not allow $w = 46 + 3.27h$
(d)	B1	for a correct numerical interpretation which includes height/h and weight/w if units are stated then they must be correct
(e)(i)	B1	awrt 546
(ii)	B1	awrt 602
(f)	B1	for 1 st estimate oe and a correct supporting reason. ignore extraneous non-contradictory comments. for 2 nd /3 rd reason, do not allow comments that refer to weights

Question Number	Scheme		Marks
2(a)	[LQ =] 95 [UQ =] 112		B1 B1 (2)
(b)	"95"−1.5×("112"−"95") or "112"+1.5×("112"−"95") "95"−1.5×("112"−"95") and "112"+1.5×("112"−"95") So only outlier is 139		M1 A1ft A1 (3)
(c)	 <p>Club A</p>		M1M1 M1A1ft (4)
(d)	(Club B has) positive (skew) So club B weights are more varied above the median (than below the median)		M1 A1 (2)
	Notes		Total 11
(a)	B1	correct lower quartile 95 cao	
	B1	correct upper quartile 112 cao	
(b)	M1	a correct method to find either the lower or upper outlier boundary ft their values from (a)	
	A1ft	correct attempt to find both the lower and upper outlier boundary ft their values from (a) this mark is for both correct ft expressions which need not be evaluated but may be implied by both of 69.5 and 137.5	
	A1	(dep on M1 only) for identifying 139 as an outlier	
(c)	M1	for a box with 1 upper whisker and 1 lower whisker	
	M1	for lower whisker plotted at 72 and upper whisker plotted at 127 or '137.5' (condone 137 to 138)'	
	M1	for quartiles plotted at 'Q ₁ ', 105 and 'Q ₃ '	
	A1ft	for a fully correct boxplot with 1 outlier marked at 139 only ft 'Q ₁ ', 'Q ₃ ' and '137.5'	
(d)	M1	for identifying positive (skew) ignore comments about the skewness of Club A positive correlation is M0	
	A1	for a correct contextual interpretation of positive skew indicating that there is greater variation above the median	

Question Number	Scheme		Marks												
3(a)	$P(Y = y) = \frac{1}{6}$ [y =] 1, 2, 3, 4, 5, 6		B1dB1 (2)												
(b)	[Discrete] uniform [distribution]		B1 (1)												
(c)	$E(Y) = \frac{6+1}{2} [= 3.5]$ or $E(Y) = \frac{1+2+3+4+5+6}{6} [= 3.5]$ $E(4Y+3) = 4 \times 3.5 + 3$		M1 M1 A1 (3)												
			Total 6												
(a)	B1	for identifying $P(Y = y) = \frac{1}{6}$ as the only probability Ignore labels. (May be seen in a table). Must be seen in part (a). Stating or using Normal, Binomial (or other distribution) is BO													
	dB1	Dep on previous B1 for $y = 1, 2, 3, 4, 5, 6$ Ignore labels. (May be seen in a table) Must be seen in part (a).													
	SC	Use of inaccurate value for $\frac{1}{6}$ can score B0B1 if otherwise fully correct e.g. <table border="1" style="display: inline-table; vertical-align: middle;"> <tr><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr> <tr><td>0.167</td><td>0.167</td><td>0.167</td><td>0.167</td><td>0.167</td><td>0.167</td></tr> </table>	1	2	3	4	5	6	0.167	0.167	0.167	0.167	0.167	0.167	
1	2	3	4	5	6										
0.167	0.167	0.167	0.167	0.167	0.167										
(b)	B1	do not allow continuous uniform [distribution] do not allow Normal uniform													
(c)	M1	a correct method to find $E(Y)$ (implied by 3.5) or sight of 7, 11, 15, 19, 23, 27 (allow one slip) $1 \times \frac{1}{6} + 2 \times \frac{2}{6} + 3 \times \frac{3}{6} + 4 \times \frac{4}{6} + 5 \times \frac{5}{6} + 6 \times \frac{6}{6}$ is MO													
	M1	use of $4E(Y) + 3$ ft their $E(Y)$ must be clear that they are using their $E(Y)$ e.g. $4\left(\frac{1}{6}\right) + 3$ is MO													

		or a correct method to find $E(4Y + 3)$ ft their 7, 11, 15, 19, 23, 27
	A1	cao 17 scores 3/3

Question Number	Scheme							Marks
4(a)	$0.05 + 0.1 + a + b + 0.35 = 1 \quad [a + b = 0.5]$ $-0.05 + a + 2b + 1.05 = 1.8 \quad [a + 2b = 0.8]$ $b = 0.3 \quad a = 0.2$							M1 M1 A1 (3)
(b)		[x]	-1	0	1	2	3	
		[F(x)]	0.0 5	0.15	0.35	0.65	1	
								B1B1ft (2)
(c)	0.15							B1 (1)
(d)	$E(X^2) = 1 \times 0.05 + 1 \times "0.2" + 4 \times "0.3" + 9 \times 0.35 [= 4.6]$ $Var(X) = "4.6" - 1.8^2 = 1.36$ $Var(5 - 3X) = 9 \times "1.36"$ $= 12.24$			$E(5 - 3X) = 8 \times 0.05 + 5 \times 0.1 + 2 \times "0.2" - 1 \times "0.3" - 4 \times 0.35 [= -0.4]$ $or \quad E(5 - 3X) = 5 - 3 \times 1.8 [= -0.4]$ $E(5 - 3X)^2 = 64 \times 0.05 + 25 \times 0.1 + 4 \times "0.2" + 1 \times "0.3" + 16 \times 0.35 [= 12.4]$ $Var(5 - 3X) = "12.4" - (" - 0.4")^2$ $= 12.24$				M1 M1 M1 A1 (4)
	Notes							Total 10

(a) **M1** for either correct equation in a and b

M1 for both correct equations in a and b

A1 both $a = 0.2$ oe and $b = 0.3$ oe
correct answers score 3 out of 3 unless from obvious incorrect working

In part (b) and (d) working labelled in script takes priority.

Look out for working shown under the table which can be marked if it does not contradict the working in the script.

(b)	B1	for 3 correct values of x and their associated cumulative probabilities
	B1ft	for a fully correct cdf ft their a and b provided both are probabilities and $a + b = 0.5$
(c)	B1	0.15 oe
(d)	M1	for method to find $E(X^2)$ with at least 3 correct (ft) non-zero products (may be implied by 4.6) only allow ft products which use probabilities or a method to find $E(5 - 3X)$ with at least 3 correct (ft) non-zero products (may be implied by -0.4) or use of $E(5 - 3X) = 5 - 3E(X)$
	M1	for a correct method to find $Var(X)$ ft their $E(X^2)$ (may be implied by 1.36) or a correct method to find $E(5 - 3X)^2$ (may be implied by 12.4)

	M1	for use of $3^2 \text{Var}(X)$ ft their $\text{Var}(X)$ or use of $E(5 - 3X)^2 - (E(5 - 3X))^2$
	A1	awrt 12.2 working must be shown Answer only send to review.

Question Number	Scheme		Marks
5(a)	Width = 7.2		B1
	Area of 21.6 for a frequency of 75 so $\frac{21.6}{75} \times 40 = 11.52$ for a frequency of 40 or $w \times h = 11.52$ or $fd = 15$		M1
	$h = \frac{11.52}{7.2} \text{ or } \frac{2}{15} \times 12 = 1.6$		A1
			(3)
(b)	$[Q_2 = 130+] \frac{5}{63} \times 10 \text{ or } [Q_2 = 130+] \frac{5.5}{63} \times 10$ or $\frac{x - 130}{140 - 130} = \frac{150 - 145}{208 - 145} \text{ or } \frac{140 - x}{140 - 130} = \frac{208 - 150}{208 - 145}$ $= \text{awrt } 130.8 \text{ or awrt } 130.9$		M1
			A1
			(2)
(c)(i)	$[\bar{y} =] \frac{6000}{300} [= 20] \text{ and } [\bar{w} =] 2 \times "20" + 200$		M1
		$= 240$	A1
			(2)
(c)(ii)	$[\text{Var}(y) =] \frac{150000}{300} - "20"^2 [= 100]$		M1
	$[\text{Var}(w) =] 4 \times "100" = 400$		M1A1
			(3)
Notes			Total 10
(a)	B1	7.2 or exact equivalent	
	M1	for sight of 11.52 or $w \times h = 11.52$ or $fd = 15$ May be implied by $h = 1.6$	
	A1	1.6 or exact equivalent	
(b)	M1	for any valid method to find the median allow use of $n+1$ allow working backwards $[140 -] \frac{58}{63} \times 10$ or $[140 -] \frac{57.5}{63} \times 10$	
	A1	awrt 130.8 allow awrt 130.9 if using $(n+1)$ No fractions	
(c)(i)	M1	for a correct method to find the mean of w Alt: $\sum w = 6000 \times 2 + 300 \times 200 [= 72000]$ and $\bar{w} = \frac{"72000"}{300}$	
	A1	cao correct answer is 2/2	
(c)(ii)	M1	for a correct method to find $\text{Var}(y)$ or for $\sum w^2 = 17400000$	
	M1	for a correct method to find $\text{Var}(w)$ ft their $\text{Var}(y)$ $2 \times "100" + 200 = 400$ is MOAO or for $\text{Var}(w) = \frac{17400000}{300} - "240"^2$	
	A1	cao answer only with no working shown scores M1MOAO	
	SC	If no marks scored in (ii), allow M1(MOAO) for	

		$\sqrt{\frac{150000}{300} - "20"^2}$
--	--	--------------------------------------

Question Number	Scheme		Marks		
6(a)	$P(A) = 0.6$	$P(B) = 0.24 + 0.09 + x$	$P(A \cap B) = 0.24$		
	$0.6(0.33 + x) = 0.24$		M1		
	$x = 0.07$		A1		
			(3)		
(b)	$P(B C') = \frac{0.24 + "0.07"}{0.6 + "0.07" + z} = \frac{31}{75}$		M1		
	$z = 0.08$		A1		
			(2)		
(c)	$y = 1 - 0.36 - 0.24 - "0.07" - 0.09 - "0.08"$		M1		
	$y = 0.16$		A1ft		
			(2)		
(d)(i) (ii)	0.36		B1		
	0.67		B1ft		
			(2)		
Notes			Total 9		
(a)	M1	for at least 2 of the 3 correct probabilities/probability expressions seen or used (implied by 2 nd M1) $0.24 + 0.09 + x$ may be implied by embedded working e.g. $x = "P(B)" - 0.24 - 0.09$			
	A1	oe correct answer scores 3/3			
(b)	M1	for a correct conditional probability equation ft "their x" substituted or just x allow $P(B C') = \frac{0.24 + "0.07"}{1 - 0.09 - y} = \frac{31}{75}$			
	A1	oe correct answer scores 2 out of 2			
(c)	M1	for a correct expression to find y ft their x and their z values substituted where x and z are probabilities do not allow blank/no answers for x and/or z as zero may be implied by $x + y + z = 0.31$ with x, y , and z as probabilities			
(d)(i) (ii)	A1ft	ft their x and their z $x + y + z = 0.31$ where x, y , and z are probabilities			
	B1	oe			
(ii)	B1ft	0.67 oe or $0.6 + "their x"$ ft "their x " where "their x " and $0.6 + "their x"$ are probabilities			

Question Number	Scheme		Marks
7(a)	$P(48.45 < W < 54.4) = P\left(Z < \frac{54.4 - 51}{1.7}\right) - P\left(Z < \frac{48.45 - 51}{1.7}\right)$ $[0.9772 - 0.0668] = 0.9104$	Cal: 0.91044... awrt 0.910	M1M1 A1 (3)
(b)	$P(53.55 < W < 54.4) = "0.9772" - (1 - "0.0668") [= 0.044]$ Calc: 0.044057... $P(W > 53.55 48.45 < W < 54.4) = \frac{"0.044"}{"0.9104"}$ $= 0.04833$	awrt 0.0483/0.0484	M1 M1 A1 (3)
(c)	$\frac{64.58 - \mu}{\sigma} = -0.5244$ (Calc: -0.524400...) $\frac{69.46 - \mu}{\sigma} = 1.2816$ (Calc: 1.281551) $4.88 = 1.806\sigma \Rightarrow \sigma = 2.702...$ awrt 2.7 $\mu = 65.996...$ awrt 66		M1 M1 dM1A1 (4)
	Notes		Total 10
(a)	M1	Standardising with 54.4, 51 and 1.7 (implied by ± 2) or standardising with 48.45, 51 and 1.7 (implied by ± 1.5) using 1.7^2 is M0	
	M1	Standardising with 54.4, 51 and 1.7 and 48.45, 51 and 1.7 and attempt to find correct probability using $p - q$ oe (where $0.5 < p < 1$ and $0 < q < 0.5$) or $p - (1 - r)$ oe (where $0.5 < p < 1$ and $0.5 < r < 1$) If 1 st M1 scored, then may be implied by correct answer	
	A1	awrt 0.910 condone 0.91 correct answer with no working scores 0	
(b)	M1	for a valid attempt to find $P(53.55 < W < 54.4)$ e.g. " $0.9772 - (1 - "0.0668")$ " May be implied by awrt 0.044/0.0441	
	M1	for a conditional probability in the form $\frac{p}{"0.9104"}$ where $0 < p < "0.9104"$ assuming independence is M0 e.g. $\frac{p \times "0.9104"}{"0.9104"}$	
	A1	awrt 0.0483/0.0484 No fractions	
(c)	M1	for standardising 64.58 and setting equal to z value where $0.5 < z < 0.6$ or for standardising 69.46 and setting equal to z value where $1 < z < 1.5$	
	M1	for standardising 64.58 and setting equal to z value where $-0.6 < z < -0.5$ oe with compatible signs and for standardising 69.46	

		and setting equal to z value where $1 < z < 1.5$ oe with compatible signs
	dM1	Dep on 1 st M1 for attempting to solve their two linear equations (using substitution or elimination) leading to an equation in one variable Implied by either $\sigma = \text{awrt } 2.7$ or $\mu = \text{awrt } 66$
	A1	both $\sigma = \text{awrt } 2.7$ and $\mu = \text{awrt } 66$ and both 0.5244 (or better) and 1.2816 (or better) seen

Question Number	Scheme		Marks
8(a)	$\frac{1}{2} \times \frac{1}{5} + \frac{1}{6} \times \frac{1}{10} + \frac{1}{3} \times \frac{1}{20} = \frac{2}{15} *$		B1* (1)
(b)	$\frac{1}{2} \times \frac{1}{5} \left[= \frac{1}{10} \right] = \frac{3}{4} *$ or $1 - \frac{1}{6} \times \frac{1}{10} + \frac{1}{3} \times \frac{1}{20} = \frac{3}{4} *$		M1A1* (2)
(c)	<pre> graph LR Root(()) -- "Late" --> Car1((Car)) Root -- "Late" --> B1((Bicycle)) Root -- "Late" --> Walk1((Walk)) Root -- "Not late" --> Car2((Car)) Root -- "Not late" --> B2((Bicycle)) Root -- "Not late" --> Walk2((Walk)) </pre>		B1B1 M1A1 (4)
	Notes		Total 7
(a)	B1*	for a correct expression which is the sum of three products of probabilities leading to the given answer allow $\frac{1}{10} + \frac{1}{60} + \frac{1}{60} = \frac{2}{15}$	
(b)	M1	for $\frac{1}{2} \times \frac{1}{5} \left[= \frac{1}{10} \right]$ (do not award if embedded in an incorrect calculation) or use of $1 - \dots$ with $\frac{1}{6} \times \frac{1}{10} + \frac{1}{3} \times \frac{1}{20}$	
	A1*	for a fully correct expression leading to the given answer	
(c)	B1	for $\frac{13}{15}$ oe in the correct place on the tree diagram	
	B1	for $\frac{1}{8}$ oe and $\frac{1}{8}$ oe in the correct place on the tree diagram	

	M1	for $\frac{4}{5} \times \frac{1}{2} \left[= \frac{6}{13} \right]$ oe or $\frac{1}{6} \times \frac{9}{10} \left[= \frac{9}{52} \right]$ oe or $\frac{1}{3} \times \frac{19}{20} \left[= \frac{19}{52} \right]$ oe implied by 1 correct probability in the correct place on tree diagram
	A1	for $\frac{6}{13}$ oe and $\frac{9}{52}$ oe and $\frac{19}{52}$ oe in the correct place on tree diagram